arXiv:1007.0613v1 [math.DS] 5 Jul 2010

Dynamics of postcritically bounded polynomial semigroups II: fiberwise dynamics and the Julia sets *

Hiroki Sumi
Department of Mathematics, Graduate School of Science
Osaka University
1-1, Machikaneyama, Toyonaka, Osaka, 560-0043, Japan
E-mail: sumi@math.sci.osaka-u.ac.jp
http://www.math.sci.osaka-u.ac.jp/∼sumi/welcomeou-e.html

Abstract

We investigate the dynamics of semigroups generated by polynomial maps on the Riemann sphere such that the postcritical set in the complex plane is bounded. Moreover, we investigate the associated random dynamics of polynomials. Furthermore, we investigate the fiberwise dynamics of skew products related to polynomial semigroups with bounded planar postcritical set. Using uniform fiberwise quasiconformal surgery on a fiber bundle, we show that if the Julia set of such a semigroup is disconnected, then there exist families of uncountably many mutually disjoint quasicircles with uniform dilatation which are parameterized by the Cantor set, densely inside the Julia set of the semigroup. Moreover, we give a sufficient condition for a fiberwise Julia set J_γ to satisfy that J_γ is a Jordan curve but not a quasicircle, the unbounded component of $\hat{\mathbb{C}} \setminus J_\gamma$ is a John domain and the bounded component of $\mathbb{C} \setminus J_\gamma$ is not a John domain. We show that under certain conditions, a random Julia set is almost surely a Jordan curve, but not a quasicircle. Many new phenomena of polynomial semigroups and random dynamics of polynomials that do not occur in the usual dynamics of polynomials are found and systematically investigated.

1 Introduction

The theory of complex dynamical systems, which has its origin in the important work of Fatou and Julia in the 1910s, has been investigated by many people and discussed in depth. In particular, since D. Sullivan showed the famous "no wandering domain theorem" using Teichmüller theory in the 1980s, this subject has attracted many researchers from a wide area. For a general reference on complex dynamical systems, see Milnor's textbook [13].

There are several areas in which we deal with generalized notions of classical iteration theory of rational functions. One of them is the theory of dynamics of rational semigroups (semigroups generated by holomorphic maps on the Riemann sphere $\hat{\mathbb{C}}$), and another one is the theory of random dynamics of holomorphic maps on the Riemann sphere.

In this paper, we will discuss these subjects. A **rational semigroup** is a semigroup generated by a family of non-constant rational maps on $\hat{\mathbb{C}}$, where $\hat{\mathbb{C}}$ denotes the Riemann sphere, with the semigroup operation being functional composition ([10]). A **polynomial semigroup** is a semigroup generated by a family of non-constant polynomial maps. Research on the dynamics of rational semigroups was initiated by A. Hinkkanen and G. J. Martin ([10]), who were interested in the role of the dynamics of polynomial semigroups while studying various one-complex-dimensional

---

*Date: July 5, 2010. 2000 Mathematics Subject Classification. 37F10, 30D05. Keywords: Polynomial semigroups, random complex dynamical systems, random iteration, skew product, Julia sets, fiberwise Julia sets.

moduli spaces for discrete groups, and by F. Ren's group([9]), who studied such semigroups from the perspective of random dynamical systems. Moreover, the research on rational semigroups is related to that on "iterated function systems" in fractal geometry. In fact, the Julia set of a rational semigroup generated by a compact family has " backward self-similarity" (cf. [20, 21]). For other research on rational semigroups, see [35, 16, 17, 33, 34], and [19]–[31].

The research on the dynamics of rational semigroups is also directly related to that on the random dynamics of holomorphic maps. The first study in this direction was by Fornaess and Sibony ([7]), and much research has followed. (See [1, 3, 4, 2, 8, 25, 28, 29, 30, 31].)

We remark that the complex dynamical systems can be used to describe some mathematical models. For example, the behavior of the population of a certain species can be described as the dynamical system of a polynomial $f(z) = az(1-z)$ such that f preserves the unit interval and the postcritical set in the plane is bounded (cf. [6]). From this point of view, it is very important to consider the random dynamics of such polynomials (see also Example 1.4). For the random dynamics of polynomials on the unit interval, see [18].

We shall give some definitions for the dynamics of rational semigroups:

Definition 1.1 ([10, 9])**.** Let G be a rational semigroup. We set

$$F(G) = \{z \in \hat{\mathbb{C}} \mid G \text{ is normal in a neighborhood of } z\},\ \ J(G) = \hat{\mathbb{C}} \setminus F(G).$$

$F(G)$ is called the **Fatou set** of G and $J(G)$ is called the **Julia set** of G. We let $\langle h_1, h_2, \ldots \rangle$ denote the rational semigroup generated by the family $\{h_i\}$. The Julia set of the semigroup generated by a single map g is denoted by $J(g)$.

Definition 1.2. For each rational map $g : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$, we set $CV(g) := \{\text{all critical values of } g : \hat{\mathbb{C}} \to \hat{\mathbb{C}}\}$. Moreover, for each polynomial map $g : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$, we set $CV^*(g) := CV(g) \setminus \{\infty\}$. For a rational semigroup G, we set

$$P(G) := \overline{\bigcup_{g \in G} CV(g)}\ (\subset \hat{\mathbb{C}}).$$

This is called the **postcritical set** of G. Furthermore, for a polynomial semigroup G, we set $P^*(G) := P(G) \setminus \{\infty\}$. This is called the **planar postcritical set** (or **finite postcritical set**) of G. We say that a polynomial semigroup G is **postcritically bounded** if $P^*(G)$ is bounded in $\mathbb{C}$.

Remark 1.3. Let G be a rational semigroup generated by a family Λ of rational maps. Then, we have that $P(G) = \overline{\cup_{g \in G \cup \{Id\}}\, g(\cup_{h \in \Lambda} CV(h))}$, where Id denotes the identity map on $\hat{\mathbb{C}}$, and that $g(P(G)) \subset P(G)$ for each $g \in G$. From this formula, one can figure out how the set $P(G)$ (resp. $P^*(G)$) spreads in $\hat{\mathbb{C}}$ (resp. $\mathbb{C}$). In fact, in Section 3.4, using the above formula, we present a way to construct examples of postcritically bounded polynomial semigroups (with some additional properties). Moreover, from the above formula, one may, in the finitely generated case, use a computer to see if a polynomial semigroup G is postcritically bounded much in the same way as one verifies the boundedness of the critical orbit for the maps $f_c(z) = z^2 + c$.

Example 1.4. Let $\Lambda := \{h(z) = cz^a(1-z)^b \mid a, b \in \mathbb{N},\ c > 0,\ c(\frac{a}{a+b})^a(\frac{b}{a+b})^b \leq 1\}$ and let G be the polynomial semigroup generated by Λ. Since for each $h \in \Lambda$, $h([0,1]) \subset [0,1]$ and $CV^*(h) \subset [0,1]$, it follows that each subsemigroup H of G is postcritically bounded.

Remark 1.5. It is well-known that for a polynomial g with $\deg(g) \geq 2$, $P^*(\langle g \rangle)$ is bounded in $\mathbb{C}$ if and only if $J(g)$ is connected ([13, Theorem 9.5]).

As mentioned in Remark 1.5, the planar postcritical set is one piece of important information regarding the dynamics of polynomials. Concerning the theory of iteration of quadratic polynomials, we have been investigating the famous "Mandelbrot set".

When investigating the dynamics of polynomial semigroups, it is natural for us to discuss the relationship between the planar postcritical set and the figure of the Julia set. The first question in this regard is: "Let G be a polynomial semigroup such that each element $g \in G$ is of degree at least two. Is $J(G)$ necessarily connected when $P^*(G)$ is bounded in $\mathbb{C}$?" The answer is **NO.** In fact, in [35, 27, 28, 17, 29, 30], we find many examples of postcritically bounded polynomial semigroups G with disconnected Julia set such that for each $g \in G$, $\deg(g) \geq 2$. Thus, it is natural to ask the following problems.

Problem 1.6. (1) Investigate what happens if $P^*(G)$ is bounded in $\mathbb{C}$ and $J(G)$ is disconnected. (2) Classify postcritically bounded polynomial semigroups.

Applying the results in [27, 28], we investigate the dynamics of every sequence, or fiberwise dynamics of the skew product associated with the generator system (cf. Section 3.1). Moreover, we investigate the random dynamics of polynomials acting on the Riemann sphere. Let us consider a polynomial semigroup G generated by a compact family Γ of polynomials. For each sequence $\gamma = (\gamma_1, \gamma_2, \gamma_3, \ldots) \in \Gamma^{\mathbb{N}}$, we examine the dynamics along the sequence γ, that is, the dynamics of the family of maps $\{\gamma_n \circ \cdots \circ \gamma_1\}_{n=1}^{\infty}$. We note that this corresponds to the fiberwise dynamics of the skew product (see Section 3.1) associated with the generator system Γ. We show that if G is postcritically bounded, $J(G)$ is disconnected, and G is generated by a compact family Γ of polynomials; then, for almost every sequence $\gamma \in \Gamma^{\mathbb{N}}$, there exists exactly one bounded component U_γ of the Fatou set of γ, the Julia set of γ has Lebesgue measure zero, there exists no non-constant limit function in U_γ for the sequence γ, and for any point $z \in U_\gamma$, the orbit along γ tends to the interior of the smallest filled-in Julia set $\hat{K}(G)$ (see Definition 2.7) of G (cf. Theorem 3.10, Corollary 3.20). Moreover, using the uniform fiberwise quasiconformal surgery ([28]), we find sub-skew products $\overline{f}$ such that $\overline{f}$ is hyperbolic (see Definition 3.9) and such that every fiberwise Julia set of $\overline{f}$ is a K-quasicircle, where K is a constant not depending on the fibers (cf. Theorem 3.10-3). Reusing the uniform fiberwise quasiconformal surgery, we show that if G is a postcritically bounded polynomial semigroup with disconnected Julia set, then for any non-empty open subset V of $J(G)$, there exists a 2-generator subsemigroup H of G such that $J(H)$ is the disjoint union of "Cantor family of quasicircles" (a family of quasicircles parameterized by a Cantor set) with uniform distortion, and such that $J(H) \cap V \neq \emptyset$ (cf. Theorem 3.13). Note that the uniform fiberwise quasiconformal surgery is based on solving uncountably many Beltrami equations (a kind of partial differential equations).

We also investigate (semi-)hyperbolic (see Definition 3.11), postcritically bounded, polynomial semigroups generated by a compact family Γ of polynomials. Let G be such a semigroup with disconnected Julia set, and suppose that there exists an element $g \in G$ such that $J(g)$ is not a Jordan curve. Then, we give a (concrete) sufficient condition for a sequence $\gamma \in \Gamma^{\mathbb{N}}$ to satisfy that the following situation $(*)$: the Julia set of γ is a Jordan curve but not a quasicircle, the basin of infinity A_γ is a John domain, and the bounded component U_γ of the Fatou set is not a John domain (cf. Theorem 3.17, Corollary 3.21). From this result, we show that for almost every sequence $\gamma \in \Gamma^{\mathbb{N}}$, situation $(*)$ holds. In fact, in this paper, under the above assumption, we find a set $\mathcal{A}$ of γ with $(*)$ which is much larger than a set $\mathcal{U}$ of γ with $(*)$ given in [28]. Moreover, we classify hyperbolic two-generator postcritically bounded polynomial semigroups G with disconnected Julia set and we also completely classify the fiberwise Julia sets J_γ in terms of the information of γ (Theorem 3.18). Note that situation $(*)$ cannot hold in the usual iteration dynamics of a single polynomial map g with $\deg(g) \geq 2$ (Remark 3.22).

The key to investigating the dynamics of postcritically bounded polynomial semigroups is the density of repelling fixed points in the Julia set ([10, 9]), which can be shown by an application of the Ahlfors five island theorem, and the lower semi-continuity of $\gamma \mapsto J_\gamma$ ([11]), which is a consequence of potential theory. Moreover, one of the keys to investigating the fiberwise dynamics of skew products is, the observation of non-constant limit functions (cf. Lemma 5.4 and [21]). The key to investigating the dynamics of semi-hyperbolic polynomial semigroups is, the continuity of the map $\gamma \mapsto J_\gamma$ (this is highly nontrivial; see [21]) and the Johnness of the basin A_γ of infinity (cf. [23]). Note that the continuity of the map $\gamma \mapsto J_\gamma$ does not hold in general, if we do not assume semi-hyperbolicity. Moreover, one of the original aspects of this paper is the idea of "combining both the theory of rational semigroups and that of random complex dynamics". It is quite natural to investigate both fields simultaneously. However, no study (except the works of the author of this paper) thus far has done so.

Furthermore, in Section 3.4 and [27, 28], we provide a way of constructing examples of postcritically bounded polynomial semigroups with some additional properties (disconnectedness of Julia set, semi-hyperbolicity, hyperbolicity, etc.) (cf. Proposition 3.23, [27, 28]). For example, by Proposition 3.23, there exists a 2-generator postcritically bounded polynomial semigroup $G = \langle h_1, h_2 \rangle$ with disconnected Julia set such that h_1 has a Siegel disk.

As we see in Example 1.4, Section 3.4, and [27, 28], it is not difficult to construct many examples, it is not difficult to verify the hypothesis "postcritically bounded", and the class of postcritically bounded polynomial semigroups is very wide.

Throughout the paper, we will see many new phenomena in polynomial semigroups or random dynamics of polynomials that do not occur in the usual dynamics of polynomials. Moreover, these new phenomena are systematically investigated.

In Section 3, we present the main results of this paper. We give some tools in Section 4. The proofs of the main results are given in Section 5.

There are many applications of the results of postcritically bounded polynomial semigroups in many directions. In the sequel [29, 25, 32], we investigate the Markov process on $\hat{\mathbb{C}}$ associated with the random dynamics of polynomials and we consider the probability $T_\infty(z)$ of tending to $\infty \in \hat{\mathbb{C}}$ starting with the initial value $z \in \hat{\mathbb{C}}$. Applying many results of [27], it will be shown in [32] that if the associated polynomial semigroup G is postcritically bounded and the Julia set is disconnected, then the chaos of the averaged system disappears due to the cooperation of generators (cooperation principle), and the function T_∞ defined on $\hat{\mathbb{C}}$ has many interesting properties which are similar to those of the Cantor function. Such a kind of "singular functions on the complex plane" appear very naturally in random dynamics of polynomials and the results of this paper (for example, the results on the space of all connected components of a Julia set) are the keys to investigating that. (The above results have been announced in [29, 25, 24, 30].)

In [27], we find many fundamental and useful results on the connected components of Julia sets of postcritically bounded polynomial semigroups. In [28], we classify (semi-)hyperbolic, postcritically bounded, compactly generated polynomial semigroups. In the sequel [17], we give some further results on postcritically bounded polynomial semigroups, by using many results in [27, 28], and this paper. Moreover, in the sequel [26], we define a new kind of cohomology theory, in order to investigate the action of finitely generated semigroups (iterated function systems), and we apply it to the study of the dynamics of postcritically bounded polynomial semigroups.

2 Preliminaries

In this section we give some basic notations and definitions, and we present some results in [27, 28], which we need to state the main results of this paper.

Definition 2.1. We set Rat : $= \{h : \hat{\mathbb{C}} \to \hat{\mathbb{C}} \mid h$ is a non-constant rational map$\}$ endowed with the topology induced by uniform convergence on $\hat{\mathbb{C}}$ with respect to the spherical distance. We set Poly $:= \{h : \hat{\mathbb{C}} \to \hat{\mathbb{C}} \mid h$ is a non-constant polynomial$\}$ endowed with the relative topology from Rat. Moreover, we set $\mathrm{Poly}_{\deg \geq 2} := \{g \in \mathrm{Poly} \mid \deg(g) \geq 2\}$ endowed with the relative topology from Rat.

Remark 2.2. Let $d \geq 1$, $\{p_n\}_{n\in\mathbb{N}}$ a sequence of polynomials of degree d, and p a polynomial. Then, $p_n \to p$ in Poly if and only if the coefficients converge appropriately and p is of degree d.

Definition 2.3. Let $\mathcal{G}$ be the set of all postcritically bounded polynomial semigroups G such that each element of G is of degree at least two. Furthermore, we set $\mathcal{G}_{con} = \{G \in \mathcal{G} \mid J(G)$ is connected$\}$ and $\mathcal{G}_{dis} = \{G \in \mathcal{G} \mid J(G)$ is disconnected$\}$.

Definition 2.4. For a polynomial semigroup G, we denote by $\mathcal{J} = \mathcal{J}_G$ the set of all connected components J of $J(G)$ such that $J \subset \mathbb{C}$. Moreover, we denote by $\hat{\mathcal{J}} = \hat{\mathcal{J}}_G$ the set of all connected components of $J(G)$.

Remark 2.5. If a polynomial semigroup G is generated by a compact set in $\mathrm{Poly}_{\deg \geq 2}$, then $\infty \in F(G)$ and thus $\mathcal{J} = \hat{\mathcal{J}}$.

Definition 2.6 ([27])**.** For any connected sets K_1 and K_2 in $\mathbb{C}$, "$K_1 \leq K_2$" indicates that $K_1 = K_2$, or K_1 is included in a bounded component of $\mathbb{C} \setminus K_2$. Furthermore, "$K_1 < K_2$" indicates $K_1 \leq K_2$ and $K_1 \neq K_2$. Note that "$\leq$" is a partial order in the space of all non-empty compact connected sets in $\mathbb{C}$. This "$\leq$" is called the **surrounding order.**

Definition 2.7 ([27])**.** For a polynomial semigroup G, we set

$$\hat{K}(G) := \{z \in \mathbb{C} \mid \bigcup_{g \in G} \{g(z)\} \text{ is bounded in } \mathbb{C}\}$$

and call $\hat{K}(G)$ the **smallest filled-in Julia set** of G. For a polynomial g, we set $K(g) := \hat{K}(\langle g \rangle)$. For a set $A \subset \hat{\mathbb{C}}$, we denote by $\mathrm{int}(A)$ the set of all interior points of A. For a polynomial semigroup G with $\infty \in F(G)$, we denote by $F_\infty(G)$ the connected component of $F(G)$ containing ∞. Moreover, for a polynomial g with $\deg(g) \geq 2$, we set $F_\infty(g) := F_\infty(\langle g \rangle)$.

The following three results in [27] are needed to state main results in this paper.

Theorem 2.8 ([27])**.** *Let $G \in \mathcal{G}$ (possibly generated by a non-compact family). Then we have all of the following.*

1. *We have $(\mathcal{J}, \leq)$ is totally ordered.*
2. *Each connected component of $F(G)$ is either simply or doubly connected.*
3. *For any $g \in G$ and any connected component J of $J(G)$, we have that $g^{-1}(J)$ is connected. Let $g^*(J)$ be the connected component of $J(G)$ containing $g^{-1}(J)$. If $J \in \mathcal{J}$, then $g^*(J) \in \mathcal{J}$. If $J_1, J_2 \in \mathcal{J}$ and $J_1 \leq J_2$, then $g^{-1}(J_1) \leq g^{-1}(J_2)$ and $g^*(J_1) \leq g^*(J_2)$.*

Theorem 2.9 ([27])**.** *Let $G \in \mathcal{G}_{dis}$ (possibly generated by a non-compact family). Then we have all of the following.*

1. *We have $\infty \in F(G)$. Thus $\mathcal{J} = \hat{\mathcal{J}}$.*
2. *The component $F_\infty(G)$ of $F(G)$ containing ∞ is simply connected. Furthermore, the element $J_{\max} = J_{\max}(G) \in \mathcal{J}$ containing $\partial F_\infty(G)$ is the unique element of $\mathcal{J}$ satisfying that $J \leq J_{\max}$ for each $J \in \mathcal{J}$.*
3. *There exists a unique element $J_{\min} = J_{\min}(G) \in \mathcal{J}$ such that $J_{\min} \leq J$ for each element $J \in \mathcal{J}$.*
4. *We have that $\mathrm{int}(\hat{K}(G)) \neq \emptyset$.*

For the figures of the Julia sets of semigroups $G \in \mathcal{G}_{dis}$, see figure 1.

Proposition 2.10 ([27])**.** *Let G be a polynomial semigroup generated by a compact subset Γ of $\mathrm{Poly}_{\deg \geq 2}$. Suppose that $G \in \mathcal{G}_{dis}$. Then, there exists an element $h_1 \in \Gamma$ with $J(h_1) \subset J_{\max}$ and there exists an element $h_2 \in \Gamma$ with $J(h_2) \subset J_{\min}$.*

3 Main results

In this section, we present the main results of this paper. The proofs of the results are given in Section 5.

3.1 Fiberwise dynamics and Julia sets

We present some results on the fiberwise dynamics of the skew product related to a postcritically bounded polynomial semigroup with disconnected Julia set. In particular, using the uniform fiberwise quasiconformal surgery on a fiber bundle, we show the existence of a family of quasicircles parameterized by a Cantor set with uniform distortion in the Julia set of such a semigroup. The proofs are given in Section 5.1.

Definition 3.1 ([21, 23])**.**

1. Let X be a compact metric space, $g : X \to X$ a continuous map, and $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ a continuous map. We say that f is a rational skew product (or fibered rational map on trivial bundle $X \times \hat{\mathbb{C}}$) over $g : X \to X$, if $\pi \circ f = g \circ \pi$ where $\pi : X \times \hat{\mathbb{C}} \to X$ denotes the canonical projection, and if for each $x \in X$, the restriction $f_x := f|_{\pi^{-1}(\{x\})} : \pi^{-1}(\{x\}) \to \pi^{-1}(\{g(x)\})$ of f is a non-constant rational map, under the canonical identification $\pi^{-1}(\{x'\}) \cong \hat{\mathbb{C}}$ for each $x' \in X$. Let $d(x) = \deg(f_x)$, for each $x \in X$. Let $f_{x,n}$ be the rational map defined by: $f_{x,n}(y) = \pi_{\hat{\mathbb{C}}}(f^n(x, y))$, for each $n \in \mathbb{N}, x \in X$ and $y \in \hat{\mathbb{C}}$, where $\pi_{\hat{\mathbb{C}}} : X \times \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ is the projection map.

   Moreover, if $f_{x,1}$ is a polynomial for each $x \in X$, then we say that $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ is a polynomial skew product over $g : X \to X$.

2. Let Γ be a compact subset of Rat. We set $\Gamma^{\mathbb{N}} := \{\gamma = (\gamma_1, \gamma_2, \ldots) \mid \forall j, \gamma_j \in \Gamma\}$ endowed with the product topology. This is a compact metric space. Let $\sigma : \Gamma^{\mathbb{N}} \to \Gamma^{\mathbb{N}}$ be the shift map, which is defined by $\sigma(\gamma_1, \gamma_2, \ldots) := (\gamma_2, \gamma_3, \ldots)$. Moreover, we define a map $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ by: $(\gamma, y) \mapsto (\sigma(\gamma), \gamma_1(y))$, where $\gamma = (\gamma_1, \gamma_2, \ldots)$. This is called **the skew product associated with the family Γ of rational maps.** Note that $f_{\gamma,n}(y) = \gamma_n \circ \cdots \circ \gamma_1(y)$.

Remark 3.2. Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a rational skew product over $g : X \to X$. Then, the function $x \mapsto d(x)$ is continuous in X.

Definition 3.3 ([21, 23])**.** Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a rational skew product over $g : X \to X$. Then, for each $x \in X$ and $n \in \mathbb{N}$, we set $f_x^n := f^n|_{\pi^{-1}(\{x\})} : \pi^{-1}(\{x\}) \to \pi^{-1}(\{g^n(x)\}) \subset X \times \hat{\mathbb{C}}$. For each $x \in X$, we denote by $F_x(f)$ the set of points $y \in \hat{\mathbb{C}}$ which has a neighborhood U in $\hat{\mathbb{C}}$ such that $\{f_{x,n} : U \to \hat{\mathbb{C}}\}_{n \in \mathbb{N}}$ is normal. Moreover, we set $F^x(f) := \{x\} \times F_x(f)$ $(\subset X \times \hat{\mathbb{C}})$. We set $J_x(f) := \hat{\mathbb{C}} \setminus F_x(f)$. Moreover, we set $J^x(f) := \{x\} \times J_x(f)$ $(\subset X \times \hat{\mathbb{C}})$. These sets $J^x(f)$ and $J_x(f)$ are called the **fiberwise Julia sets**. Moreover, we set $\tilde{J}(f) := \overline{\bigcup_{x \in X} J^x(f)}$, where the closure is taken in the product space $X \times \hat{\mathbb{C}}$. For each $x \in X$, we set $\hat{J}^x(f) := \pi^{-1}(\{x\}) \cap \tilde{J}(f)$. Moreover, we set $\hat{J}_x(f) := \pi_{\hat{\mathbb{C}}}(\hat{J}^x(f))$. We set $\tilde{F}(f) := (X \times \hat{\mathbb{C}}) \setminus \tilde{J}(f)$.

Remark 3.4. We have $\hat{J}^x(f) \supset J^x(f)$ and $\hat{J}_x(f) \supset J_x(f)$. However, strict containment can occur. For example, let h_1 be a polynomial having a Siegel disk with center $z_1 \in \mathbb{C}$. Let h_2 be a polynomial such that z_1 is a repelling fixed point of h_2. Let $\Gamma = \{h_1, h_2\}$. Let $f : \Gamma \times \hat{\mathbb{C}} \to \Gamma \times \hat{\mathbb{C}}$ be the skew product associated with the family Γ. Let $x = (h_1, h_1, h_1, \ldots) \in \Gamma^{\mathbb{N}}$. Then, $(x, z_1) \in \hat{J}^x(f) \setminus J^x(f)$ and $z_1 \in \hat{J}_x(f) \setminus J_x(f)$.

Definition 3.5. Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a polynomial skew product over $g : X \to X$. Then for each $x \in X$, we set $K_x(f) := \{y \in \hat{\mathbb{C}} \mid \{f_{x,n}(y)\}_{n \in \mathbb{N}} \text{ is bounded in } \mathbb{C}\}$, and $A_x(f) := \{y \in \hat{\mathbb{C}} \mid f_{x,n}(y) \to \infty,\ n \to \infty\}$. Moreover, we set $K^x(f) := \{x\} \times K_x(f)$ $(\subset X \times \hat{\mathbb{C}})$ and $A^x(f) := \{x\} \times A_x(f)$ $(\subset X \times \hat{\mathbb{C}})$.

Definition 3.6. Let G be a polynomial semigroup generated by a subset Γ of $\text{Poly}_{\deg \geq 2}$. Suppose $G \in \mathcal{G}_{dis}$. Then we set $\Gamma_{\min} := \{h \in \Gamma \mid J(h) \subset J_{\min}\}$, where $J_{\min}$ denotes the unique minimal element in $(\mathcal{J}, \leq)$ in Theorem 2.9-3. Furthermore, if $\Gamma_{\min} \neq \emptyset$, let $G_{\min,\Gamma}$ be the subsemigroup of G that is generated by $\Gamma_{\min}$.

Remark 3.7. Let G be a polynomial semigroup generated by a compact subset Γ of $\text{Poly}_{\deg\geq 2}$. Suppose $G \in \mathcal{G}_{dis}$. Then, by Proposition 2.10, we have $\Gamma_{\min} \neq \emptyset$ and $\Gamma \setminus \Gamma_{\min} \neq \emptyset$. Moreover, $\Gamma_{\min}$ is a compact subset of Γ. For, if $\{h_n\}_{n\in\mathbb{N}} \subset \Gamma_{\min}$ and $h_n \to h_\infty$ in Γ, then for a repelling periodic point $z_0 \in J(h_\infty)$ of h_∞, we have that $d(z_0, J(h_n)) \to 0$ as $n \to \infty$, which implies that $z_0 \in J_{\min}$ and thus $h_\infty \in \Gamma_{\min}$.

Notation: Let $\mathcal{F} := \{\varphi_n\}_{n\in\mathbb{N}}$ be a sequence of meromorphic functions in a domain V. We say that a meromorphic function ψ is a limit function of $\mathcal{F}$ if there exists a strictly increasing sequence $\{n_j\}_{j\in\mathbb{N}}$ of positive integers such that $\varphi_{n_j} \to \psi$ locally uniformly on V, as $j \to \infty$.

Definition 3.8. Let Γ and S be non-empty subsets of $\text{Poly}_{\deg\geq 2}$ with $S \subset \Gamma$. We set

$$R(\Gamma, S) := \left\{\gamma = (\gamma_1, \gamma_2, \ldots) \in \Gamma^{\mathbb{N}} \mid \sharp(\{n \in \mathbb{N} \mid \gamma_n \in S\}) = \infty\right\}.$$

Definition 3.9. Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a rational skew product over $g : X \to X$. We set

$$C(f) := \{(x, y) \in X \times \hat{\mathbb{C}} \mid y \text{ is a critical point of } f_{x,1}\}.$$

Moreover, we set $P(f) := \overline{\cup_{n\in\mathbb{N}} f^n(C(f))}$, where the closure is taken in the product space $X \times \hat{\mathbb{C}}$. This $P(f)$ is called the **fiber-postcritical set** of f.

We say that f is hyperbolic (along fibers) if $P(f) \subset F(f)$.

We present a result which describes the details of the fiberwise dynamics along γ in $R(\Gamma, \Gamma \setminus \Gamma_{\min})$.

Theorem 3.10. *Let G be a polynomial semigroup generated by a compact subset Γ of $\text{Poly}_{\deg\geq 2}$. Suppose $G \in \mathcal{G}_{dis}$. Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with the family Γ of polynomials. Then, all of the following statements 1,2, and 3 hold.*

1. *Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$. Then, each limit function of $\{f_{\gamma,n}\}_{n\in\mathbb{N}}$ in each connected component of $F_\gamma(f)$ is constant.*
2. *Let S be a non-empty compact subset of $\Gamma \setminus \Gamma_{\min}$. Then, for each $\gamma \in R(\Gamma, S)$, we have the following.*
    - (a) *There exists exactly one bounded component U_γ of $F_\gamma(f)$. Furthermore, $\partial U_\gamma = \partial A_\gamma(f) = J_\gamma(f)$.*
    - (b) *For each $y \in U_\gamma$, there exists a number $n \in \mathbb{N}$ such that $f_{\gamma,n}(y) \in \text{int}(\hat{K}(G))$.*
    - (c) *$\hat{J}_\gamma(f) = J_\gamma(f)$. Moreover, the map $\omega \mapsto J_\omega(f)$ defined on $\Gamma^{\mathbb{N}}$ is continuous at γ, with respect to the Hausdorff metric in the space of non-empty compact subsets of $\hat{\mathbb{C}}$.*
    - (d) *The 2-dimensional Lebesgue measure of $\hat{J}_\gamma(f) = J_\gamma(f)$ is equal to zero.*
3. *Let S be a non-empty compact subset of $\Gamma \setminus \Gamma_{\min}$. For each $p \in \mathbb{N}$, we denote by $W_{S,p}$ the set of elements $\gamma = (\gamma_1, \gamma_2, \ldots) \in \Gamma^{\mathbb{N}}$ such that for each $l \in \mathbb{N}$, at least one of $\gamma_{l+1}, \ldots, \gamma_{l+p}$ belongs to S. Let $\overline{f} := f|_{W_{S,p}\times\hat{\mathbb{C}}} : W_{S,p} \times \hat{\mathbb{C}} \to W_{S,p} \times \hat{\mathbb{C}}$. Then, $\overline{f}$ is a hyperbolic skew product over the shift map $\sigma : W_{S,p} \to W_{S,p}$, and there exists a constant $K_{S,p} \geq 1$ such that for each $\gamma \in W_{S,p}$, $\hat{J}_\gamma(f) = J_\gamma(f) = J_\gamma(\overline{f})$ is a $K_{S,p}$-quasicircle. Here, a Jordan curve ξ in $\hat{\mathbb{C}}$ is said to be a K-quasicircle, if ξ is the image of $S^1 (\subset \mathbb{C})$ under a K-quasiconformal homeomorphism $\varphi : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$. (For the definition of a quasicircle and a quasiconformal homeomorphism, see [12].)*

Definition 3.11. Let G be a rational semigroup.

1. We say that G is hyperbolic if $P(G) \subset F(G)$.

2. We say that G is semi-hyperbolic if there exists a number $\delta > 0$ and a number $N \in \mathbb{N}$ such that for each $y \in J(G)$ and each $g \in G$, we have $\deg(g : V \to B(y, \delta)) \leq N$ for each connected component V of $g^{-1}(B(y, \delta))$, where $B(y, \delta)$ denotes the ball of radius δ with center y with respect to the spherical distance, and $\deg(g : \cdot \to \cdot)$ denotes the degree of finite branched covering. (For background of semi-hyperbolicity, see [21] and [23].)

Theorem 3.12. *Let G be a polynomial semigroup generated by a compact subset Γ of $\text{Poly}_{\deg \geq 2}$. Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with the family Γ. Suppose that $G \in \mathcal{G}_{dis}$ and that G is semi-hyperbolic. Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$ be any element. Then, $\hat{J}_\gamma(f) = J_\gamma(f)$ and $J_\gamma(f)$ is a Jordan curve. Moreover, for each point $y_0 \in \text{int}(K_\gamma(f))$, there exists an $n \in \mathbb{N}$ such that $f_{\gamma,n}(y_0) \in \text{int}(\hat{K}(G))$.*

We next present a result that there exist families of uncountably many mutually disjoint quasicircles with uniform distortion, densely inside the Julia set of a semigroup in $\mathcal{G}_{dis}$.

Theorem 3.13. (Existence of a Cantor family of quasicircles.) *Let $G \in \mathcal{G}_{dis}$ (possibly generated by a non-compact family) and let V be an open subset of $\hat{\mathbb{C}}$ with $V \cap J(G) \neq \emptyset$. Then, there exist elements g_1 and g_2 in G such that all of the following hold.*

1. *$H = \langle g_1, g_2 \rangle$ satisfies that $J(H) \subset J(G)$.*
2. *There exists a non-empty open set U in $\hat{\mathbb{C}}$ such that $g_1^{-1}(\overline{U}) \cup g_2^{-1}(\overline{U}) \subset U$, and such that $g_1^{-1}(\overline{U}) \cap g_2^{-1}(\overline{U}) = \emptyset$.*
3. *$H = \langle g_1, g_2 \rangle$ is a hyperbolic polynomial semigroup.*
4. *Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with the family $\Gamma = \{g_1, g_2\}$ of polynomials. Then, we have the following.*
    (a) *$J(H) = \bigcup_{\gamma \in \Gamma^{\mathbb{N}}} J_\gamma(f)$ (disjoint union).*
    (b) *For each connected component J of $J(H)$, there exists an element $\gamma \in \Gamma^{\mathbb{N}}$ such that $J = J_\gamma(f)$.*
    (c) *There exists a constant $K \geq 1$ independent of J such that each connected component J of $J(H)$ is a K-quasicircle.*
    (d) *The map $\gamma \mapsto J_\gamma(f)$, defined for all $\gamma \in \Gamma^{\mathbb{N}}$, is continuous with respect to the Hausdorff metric in the space of non-empty compact subsets of $\hat{\mathbb{C}}$, and injective.*
    (e) *For each element $\gamma \in \Gamma^{\mathbb{N}}$, $J_\gamma(f) \cap V \neq \emptyset$.*
    (f) *Let $\omega \in \Gamma^{\mathbb{N}}$ be an element such that $\sharp(\{j \in \mathbb{N} \mid \omega_j = g_1\}) = \infty$ and such that $\sharp(\{j \in \mathbb{N} \mid \omega_j = g_2\}) = \infty$. Then, $J_\omega(f)$ does not meet the boundary of any connected component of $F(G)$.*

Remark 3.14. This "Cantor family of quasicircles" in the research of rational semigroups was introduced by the author of this paper. By using this idea, in [17] (which was written after this paper), it is shown that for a polynomial semigroup $G \in \mathcal{G}_{dis}$ which is generated by a (possibly non-compact) family of $\text{Poly}_{\deg \geq 2}$, if A and B are two different doubly connected components of $F(G)$, then there exists a Cantor family $\mathcal{C}$ of quasicircles in $J(G)$ such that each element of $\mathcal{C}$ separates $\overline{A}$ and $\overline{B}$.

3.2 Fiberwise Julia sets that are Jordan curves but not quasicircles

We present a result on a sufficient condition for a fiberwise Julia set $J_x(f)$ to satisfy that $J_x(f)$ is a Jordan curve but not a quasicircle, the unbounded component of $\hat{\mathbb{C}} \setminus J_x(f)$ is a John domain, and the bounded component of $\mathbb{C} \setminus J_x(f)$ is not a John domain. Note that we have many examples of this phenomenon (see Proposition 3.23,Remark 3.24,Example 3.25), and note also that this phenomenon cannot hold in the usual iteration dynamics of a single polynomial map g with $\deg(g) \geq 2$ (see Remark 3.22). The proofs are given in Section 5.2.

Definition 3.15. Let V be a subdomain of $\hat{\mathbb{C}}$ such that $\partial V \subset \mathbb{C}$. We say that V is a John domain if there exists a constant $c > 0$ and a point $z_0 \in V$ ($z_0 = \infty$ when $\infty \in V$) satisfying the following: for all $z_1 \in V$ there exists an arc $\xi \subset V$ connecting z_1 to z_0 such that for any $z \in \xi$, we have $\min\{|z-a| \mid a \in \partial V\} \geq c|z - z_1|$.

Remark 3.16. Let V be a simply connected domain in $\hat{\mathbb{C}}$ such that $\partial V \subset \mathbb{C}$. It is well-known that if V is a John domain, then ∂V is locally connected ([14, page 26]). Moreover, a Jordan curve $\xi \subset \mathbb{C}$ is a quasicircle if and only if both components of $\hat{\mathbb{C}} \setminus \xi$ are John domains ([14, Theorem 9.3]).

Theorem 3.17. *Let G be a polynomial semigroup generated by a compact subset Γ of $\mathrm{Poly}_{\deg \geq 2}$. Suppose that $G \in \mathcal{G}_{dis}$. Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with the family Γ of polynomials. Let $m \in \mathbb{N}$ and suppose that there exists an element $(h_1, h_2, \ldots, h_m) \in \Gamma^m$ such that $J(h_m \circ \cdots \circ h_1)$ is not a quasicircle. Let $\alpha = (\alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$ be the element such that for each $k, l \in \mathbb{N} \cup \{0\}$ with $1 \leq l \leq m$, $\alpha_{km+l} = h_l$. Then, the following statements 1 and 2 hold.*

1. *Suppose that G is hyperbolic. Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$ be an element such that there exists a sequence $\{n_k\}_{k\in\mathbb{N}}$ of positive integers satisfying that $\sigma^{n_k}(\gamma) \to \alpha$ as $k \to \infty$. Then, $J_\gamma(f)$ is a Jordan curve but not a quasicircle. Moreover, the unbounded component $A_\gamma(f)$ of $F_\gamma(f)$ is a John domain, but the unique bounded component U_γ of $F_\gamma(f)$ is not a John domain.*
2. *Suppose that G is semi-hyperbolic. Let $\rho_0 \in \Gamma \setminus \Gamma_{\min}$ be any element and let $\beta := (\rho_0, \alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$. Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$ be an element such that there exists a sequence $\{n_k\}_{k\in\mathbb{N}}$ of positive integers satisfying that $\sigma^{n_k}(\gamma) \to \beta$ as $k \to \infty$. Then, $J_\gamma(f)$ is a Jordan curve but not a quasicircle. Moreover, the unbounded component $A_\gamma(f)$ of $F_\gamma(f)$ is a John domain, but the unique bounded component U_γ of $F_\gamma(f)$ is not a John domain.*

We now classify hyperbolic two-generator polynomial semigroups in $\mathcal{G}_{dis}$. Moreover, we completely classify the fiberwise Julia sets $J_\gamma(f)$ in terms of the information on γ.

Theorem 3.18. *Let $\Gamma = \{h_1, h_2\} \subset \mathrm{Poly}_{\deg \geq 2}$. Let $G = \langle h_1, h_2 \rangle$. Suppose that $G \in \mathcal{G}_{dis}$ and G is hyperbolic. Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with Γ. Then, for each connected component J of $J(G)$, there exists a unique $\gamma \in \Gamma^{\mathbb{N}}$ such that $J = J_\gamma(f)$. Moreover, exactly one of the following statements 1, 2 holds.*

1. *There exists a constant $K \geq 1$ such that for each $\gamma \in \Gamma^{\mathbb{N}}$, $J_\gamma(f)$ is a K-quasicircle.*
2. *There exists a $j \in \{1, 2\}$ such that $J(h_j)$ is not a Jordan curve. In this case, for each $\gamma = (\gamma_1, \gamma_2, \ldots) \in \Gamma^{\mathbb{N}}$, exactly one of the following statements* (a),(b), (c) *holds.*
   (a) *There exists a $p \in \mathbb{N}$ such that for each $l \in \mathbb{N}$, at least one of $\gamma_{l+1}, \ldots, \gamma_{l+p}$ is not equal to h_j. Moreover, $J_\gamma(f)$ is a quasicircle.*
   (b) *$\sharp\{n \in \mathbb{N} \mid \gamma_n \neq h_j\} = \infty$ and there exists a strictly increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ in $\mathbb{N}$ such that $\sigma^{n_k}(\gamma) \to (h_j, h_j, h_j, \ldots)$ as $k \to \infty$. Moreover, $J_\gamma(f)$ is a Jordan curve but not a quasicircle, the unbounded component $A_\gamma(f)$ of $\hat{\mathbb{C}} \setminus J_\gamma(f)$ is a John domain, and the bounded component of $\mathbb{C} \setminus J_\gamma(f)$ is not a John domain.*
   (c) *There exists an $l \in \mathbb{N}$ such that $\sigma^l(\gamma) = (h_j, h_j, h_j, \ldots)$. Moreover, $J_\gamma(f)$ is not a Jordan curve.*

3.3 Random dynamics of polynomials

In this section, we present some results on the random dynamics of polynomials. The proofs are given in Section 5.3.

Let τ be a Borel probability measure on $\text{Poly}_{\deg\geq 2}$. We consider the i.i.d. random dynamics on $\hat{\mathbb{C}}$ such that at every step we choose a polynomial map $h : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ according to the distribution τ. (Hence, this is a kind of Markov process on $\hat{\mathbb{C}}$.)

Notation: For a Borel probability measure τ on $\text{Poly}_{\deg\geq 2}$, we denote by Γ_τ the support of τ on $\text{Poly}_{\deg\geq 2}$. (Hence, Γ_τ is a closed set in $\text{Poly}_{\deg\geq 2}$.) Moreover, we denote by $\tilde{\tau}$ the infinite product measure $\otimes_{j=1}^{\infty}\tau$. This is a Borel probability measure on $\Gamma_\tau^{\mathbb{N}}$.

Definition 3.19. Let X be a complete metric space. A subset A of X is said to be residual if $X \setminus A$ is a countable union of nowhere dense subsets of X. Note that by Baire Category Theorem, a residual set A is dense in X.

Corollary 3.20. *(Corollary of Theorem 3.10-2) Let* Γ *be a non-empty compact subset of* $\text{Poly}_{\deg\geq 2}$. *Let* $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ *be the skew product associated with the family* Γ *of polynomials. Let* G *be the polynomial semigroup generated by* Γ. *Suppose* $G \in \mathcal{G}_{dis}$. *Then, there exists a residual subset* $\mathcal{U}$ *of* $\Gamma^{\mathbb{N}}$ *such that for each Borel probability measure* τ *on* $\text{Poly}_{\deg\geq 2}$ *with* $\Gamma_\tau = \Gamma$, *we have* $\tilde{\tau}(\mathcal{U}) = 1$, *and such that each* $\gamma \in \mathcal{U}$ *satisfies all of the following.*

1. *There exists exactly one bounded component* U_γ *of* $F_\gamma(f)$. *Furthermore,* $\partial U_\gamma = \partial A_\gamma(f) = J_\gamma(f)$.
2. *Each limit function of* $\{f_{\gamma,n}\}_n$ *in* U_γ *is constant. Moreover, for each* $y \in U_\gamma$, *there exists a number* $n \in \mathbb{N}$ *such that* $f_{\gamma,n}(y) \in \text{int}(\hat{K}(G))$.
3. $\hat{J}_\gamma(f) = J_\gamma(f)$. *Moreover, the map* $\omega \mapsto J_\omega(f)$ *defined on* $\Gamma^{\mathbb{N}}$ *is continuous at* γ, *with respect to the Hausdorff metric in the space of non-empty compact subsets of* $\hat{\mathbb{C}}$.
4. *The 2-dimensional Lebesgue measure of* $\hat{J}_\gamma(f) = J_\gamma(f)$ *is equal to zero.*

Corollary 3.21. *(Corollary of Theorems 3.12, 3.17) Let* Γ *be a non-empty compact subset of* $\text{Poly}_{\deg\geq 2}$. *Let* $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ *be the skew product associated with the family* Γ *of polynomials. Let* G *be the polynomial semigroup generated by* Γ. *Suppose* $G \in \mathcal{G}_{dis}$ *and* G *is semi-hyperbolic. Then, we have both of the following.*

1. *There exists a residual subset* $\mathcal{U}$ *of* $\Gamma^{\mathbb{N}}$ *such that for each Borel probability measure* τ *on* $\text{Poly}_{\deg\geq 2}$ *with* $\Gamma_\tau = \Gamma$, *we have* $\tilde{\tau}(\mathcal{U}) = 1$, *and such that for each* $\gamma \in \mathcal{U}$ *and for each point* $y_0 \in int(K_\gamma(f))$, J_γ *is a Jordan curve and there exist an* $n \in \mathbb{N}$ *with* $f_{\gamma,n}(y_0) \in int(\hat{K}(G))$.
2. *Suppose further that there exists an element* $h \in G$ *such that* $J(h)$ *is not a quasicircle. Then, there exists a residual subset* $\mathcal{V}$ *of* $\Gamma^{\mathbb{N}}$ *such that for each Borel probability measure* τ *on* $\text{Poly}_{\deg\geq 2}$ *with* $\Gamma_\tau = \Gamma$, *we have* $\tilde{\tau}(\mathcal{V}) = 1$, *and such that for each* $\gamma \in \mathcal{V}$, J_γ *is a Jordan curve but not a quasicircle, the unbounded component of* $\hat{\mathbb{C}} \setminus J_\gamma$ *is a John domain, and the bounded component of* $\mathbb{C} \setminus J_\gamma$ *is not a John domain.*

Remark 3.22. Let $h \in \text{Poly}_{\deg\geq 2}$ be a polynomial. Suppose that $J(h)$ is a Jordan curve but not a quasicircle. Then, it is easy to see that there exists a parabolic fixed point of h in $\mathbb{C}$ and the bounded connected component of $F(h)$ is the immediate parabolic basin. Hence, $\langle h \rangle$ is not semi-hyperbolic. Moreover, by [5], $F_\infty(h)$ is not a John domain.

Thus what we see in Theorem 3.17 and statement 2 in Corollary 3.21, as illustrated in Example 3.25, is a special and new phenomenon which can hold in the *random* dynamics of a family of polynomials, but cannot hold in the usual iteration dynamics of a single polynomial. Namely, it can hold that for almost every $\gamma \in \Gamma^{\mathbb{N}}$, $J_\gamma(f)$ is a Jordan curve and fails to be a quasicircle all while the basin of infinity $A_\gamma(f)$ is still a John domain. Whereas, if $J(h)$, for some polynomial h, is a Jordan curve which fails to be a quasicircle, then the basin of infinity $F_\infty(h)$ is necessarily **not** a John domain.

Pilgrim and Tan Lei ([15]) showed that there exists a hyperbolic rational map h with *disconnected* Julia set such that "almost every" connected component of $J(h)$ is a Jordan curve but not a quasicircle.

3.4 Examples

We give some examples of semigroups G in $\mathcal{G}_{dis}$. The following proposition was proved in [27].

Proposition 3.23 ([27]). *Let G be a polynomial semigroup generated by a compact subset Γ of $\mathrm{Poly}_{\deg\geq 2}$. Suppose that $G \in \mathcal{G}$ and $\mathrm{int}(\hat{K}(G)) \neq \emptyset$. Let $b \in \mathrm{int}(\hat{K}(G))$. Moreover, let $d \in \mathbb{N}$ be any positive integer such that $d \geq 2$, and such that $(d, \deg(h)) \neq (2,2)$ for each $h \in \Gamma$. Then, there exists a number $c > 0$ such that for each $a \in \mathbb{C}$ with $0 < |a| < c$, there exists a compact neighborhood V of $g_a(z) = a(z-b)^d + b$ in $\mathrm{Poly}_{\deg\geq 2}$ satisfying that for any non-empty subset V' of V, the polynomial semigroup $H_{\Gamma,V'}$ generated by the family $\Gamma \cup V'$ belongs to $\mathcal{G}_{dis}$, $\hat{K}(H_{\Gamma,V'}) = \hat{K}(G)$ and $(\Gamma \cup V')_{\min} \subset \Gamma$. Moreover, in addition to the assumption above, if G is semi-hyperbolic (resp. hyperbolic), then the above $H_{\Gamma,V'}$ is semi-hyperbolic (resp. hyperbolic).*

Remark 3.24. By Proposition 3.23, there exists a 2-generator polynomial semigroup $G = \langle h_1, h_2 \rangle$ in $\mathcal{G}_{dis}$ such that h_1 has a Siegel disk. Moreover, by Proposition 3.23, we can easily construct many examples G that satisfies statements 1, 2 in Theorem 3.17 and statement 2 in Corollary 3.21.

Example 3.25. Let $g_1(z) := z^2 - 1$ and $g_2(z) := \frac{z^2}{4}$. Let $\Gamma := \{g_1^2, g_2^2\}$. Moreover, let G be the polynomial semigroup generated by Γ. Let $D := \{z \in \mathbb{C} \mid |z| < 0.4\}$. Then, it is easy to see $g_1^2(D) \cup g_2^2(D) \subset D$. Hence, $D \subset F(G)$. Moreover, by Remark 1.3, we have that $P^*(G) = \cup_{g \in G \cup \{Id\}} g(\{0, -1\}) \subset D \subset F(G)$. Hence, $G \in \mathcal{G}$ and G is hyperbolic. Furthermore, let $K := \{z \in \mathbb{C} \mid 0.4 \leq |z| \leq 4\}$. Then, it is easy to see that $(g_1^2)^{-1}(K) \cup (g_2^2)^{-1}(K) \subset K$ and $(g_1^2)^{-1}(K) \cap (g_2^2)^{-1}(K) = \emptyset$. Combining it with [10, Corollary 3.2] and [20, Lemma 2.4], we obtain that $J(G)$ is disconnected. Therefore, $G \in \mathcal{G}_{dis}$. Moreover, it is easy to see that $\Gamma_{\min} = \{g_1^2\}$. Since $J(g_1^2)$ is not a Jordan curve, we can apply Theorem 3.17. Setting $\alpha := (g_1^2, g_1^2, g_1^2, \ldots) \in \Gamma^{\mathbb{N}}$, it follows that for any

$$\gamma \in \mathcal{I} := \{\omega \in R(\Gamma, \Gamma \setminus \Gamma_{\min}) \mid \exists (n_k) \text{ with } \sigma^{n_k}(\omega) \to \alpha\},$$

$J_\gamma(f)$ is a Jordan curve but not a quasicircle, and $A_\gamma(f)$ is a John domain but the bounded component of $F_\gamma(f)$ is not a John domain. (See figure 1: the Julia set of G. In this example, we have $(g_1^2)^{-1}(J(G)) \cap (g_2^2)^{-1}(J(G)) = \emptyset$, and so $\hat{\mathcal{J}}_G = \{J_\gamma(f) \mid \gamma \in \Gamma^{\mathbb{N}}\}$, and if $\gamma \neq \omega$, $J_\gamma(f) \cap J_\omega(f) = \emptyset$.) Note that by Theorem 3.18, if $\gamma \notin \mathcal{I}$, then either $J_\gamma(f)$ is not a Jordan curve or $J_\gamma(f)$ is a quasicircle.

Figure 1: The Julia set of $G = \langle g_1^2, g_2^2 \rangle$.



In [27, 28, 26, 17, 33, 29], we find many examples of postcritically bounded polynomial semigroups with many additional properties. In fact, several systematic ways to give such examples are found in those papers.

4 Tools

In this section, we introduce some fundamental tools to prove the main results.

Let G be a rational semigroup. Then, for each $g \in G$, $g(F(G)) \subset F(G), g^{-1}(J(G)) \subset J(G)$. If G is generated by a compact family Λ of Rat, then $J(G) = \bigcup_{h \in \Lambda} h^{-1}(J(G))$ (this is called the backward self-similarity). If $\sharp J(G) \geq 3$, then $J(G)$ is a perfect set and $J(G)$ is equal to the closure of the set of repelling cycles of elements of G. We set $E(G) := \{z \in \hat{\mathbb{C}} \mid \sharp \bigcup_{g \in G} g^{-1}(\{z\}) < \infty\}$. If $\sharp J(G) \geq 3$, then $\sharp E(G) \leq 2$ and for each $z \in J(G) \setminus E(G)$, $J(G) = \overline{\bigcup_{g \in G} g^{-1}(\{z\})}$. If $\sharp J(G) \geq 3$, then $J(G)$ is the smallest set in $\{\emptyset \neq K \subset \hat{\mathbb{C}} \mid K \text{ is compact}, \forall g \in G, g(K) \subset K\}$. For more details on these properties of rational semigroups, see [10, 9, 20]. For the dynamics of postcritically bounded polynomial semigroups, see [27, 28, 17]. For some fundamental properties of skew products, see [21, 28].

5 Proofs

In this section, we give the proofs of the main results.

5.1 Proofs of results in 3.1

In this section, we prove results in section 3.1.

To prove results in 3.1, we need the following notations and lemmas.

Definition 5.1 ([21]). Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a rational skew product over $g : X \to X$. Let $N \in \mathbb{N}$. We say that a point $(x_0, y_0) \in X \times \hat{\mathbb{C}}$ belongs to $SH_N(f)$ if there exists a neighborhood U of x_0 in X and a positive number δ such that for any $x \in U$, any $n \in \mathbb{N}$, any $x_n \in g^{-n}(x)$, and any connected component V of $(f_{x_n,n})^{-1}(B(y_0, \delta))$, $\deg(f_{x_n,n} : V \to B(y_0, \delta)) \leq N$. Moreover, we set $UH(f) := (X \times \hat{\mathbb{C}}) \setminus \cup_{N \in \mathbb{N}} SH_N(f)$. We say that f is semi-hyperbolic (along fibers) if $UH(f) \subset \tilde{F}(f)$.

Remark 5.2. Under the above notation, we have $UH(f) \subset P(f)$.

Remark 5.3. Let Γ be a compact subset of Rat and let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with Γ. Let G be the rational semigroup generated by Γ. Then, by [28, Remark 2.12], f is semi-hyperbolic if and only if G is semi-hyperbolic. Similarly, f is hyperbolic if and only if G is hyperbolic.

Lemma 5.4. *Let $f : X \times \hat{\mathbb{C}} \to X \times \hat{\mathbb{C}}$ be a polynomial skew product over $g : X \to X$ such that for each $\omega \in X$, $d(\omega) \geq 2$. Let $x \in X$ be a point and $y_0 \in F_x(f)$ a point. Suppose that there exists a strictly increasing sequence $\{n_j\}_{j \in \mathbb{N}}$ of positive integers such that the sequence $\{f_{x,n_j}\}_{j \in \mathbb{N}}$ converges to a non-constant map around y_0, and such that $\lim_{j \to \infty} f^{n_j}(x, y_0)$ exists. We set $(x_\infty, y_\infty) := \lim_{j \to \infty} f^{n_j}(x, y_0)$. Then, there exists a non-empty bounded open set V in $\mathbb{C}$ and a number $k \in \mathbb{N}$ such that $\{x_\infty\} \times \partial V \subset \tilde{J}(f) \cap UH(f) \subset \tilde{J}(f) \cap P(f)$, and such that for each j with $j \geq k$, $f_{x,n_j}(y_0) \in V$.*

Proof. We set

$$V := \{y \in \hat{\mathbb{C}} \mid \exists \epsilon > 0, \lim_{i \to \infty} \sup_{j > i} \sup_{d(\xi, y) \leq \epsilon} d(f_{g^{n_i}(x), n_j - n_i}(\xi), \xi) = 0\}.$$

Then, by [21, Lemma 2.13], we have $\{x_\infty\} \times \partial V \subset \tilde{J}(f) \cap UH(f) \subset \tilde{J}(f) \cap P(f)$. Moreover, since for each $x \in X$, $f_{x,1}$ is a polynomial with $d(x) \geq 2$, [28, Lemma 3.4.4] implies that there exists a ball B around ∞ such that $B \subset \hat{\mathbb{C}} \setminus V$.

From the assumption, there exists a number $a > 0$ and a non-constant map $\varphi : D(y_0, a) \to \hat{\mathbb{C}}$ such that $f_{x,n_j} \to \varphi$ as $j \to \infty$, uniformly on $D(y_0, a)$. Hence, $d(f_{x,n_j}(y), f_{x,n_i}(y)) \to 0$ as $i, j \to \infty$, uniformly on $D(y_0, a)$. Moreover, since φ is not constant, there exists a positive number ϵ such that for each large i, $f_{x,n_i}(D(y_0, a)) \supset D(y_\infty, \epsilon)$. Therefore, it follows that $d(f_{g^{n_i}(x), n_j - n_i}(\xi), \xi) \to 0$ as $i, j \to \infty$ uniformly on $D(y_\infty, \epsilon)$. Thus, $y_\infty \in V$. Hence, there exists a number $k \in \mathbb{N}$ such that for each $j \geq k$, $f_{x,n_j}(y_0) \in V$. Therefore, we have proved Lemma 5.4. □

Remark 5.5. In [21, Lemma 2.13] and [23, Theorem 2.6], the sequence (n_j) of positive integers should be strictly increasing.

Lemma 5.6. *Let* Γ *be a non-empty compact subset of* $\text{Poly}_{\deg \geq 2}$. *Let* $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ *be the skew product associated with* Γ. *Let* G *be the polynomial semigroup generated by* Γ. *Let* $\gamma \in \Gamma^{\mathbb{N}}$ *be a point. Let* $y_0 \in F_\gamma(f)$ *and suppose that there exists a strictly increasing sequence* $\{n_j\}_{j \in \mathbb{N}}$ *of positive integers such that* $\{f_{\gamma,n_j}\}_{j \in \mathbb{N}}$ *converges to a non-constant map around* y_0. *Moreover, suppose that* $G \in \mathcal{G}$. *Then, there exists a number* $j \in \mathbb{N}$ *such that* $f_{\gamma,n_j}(y_0) \in \text{int}(\hat{K}(G))$.

Proof. By Lemma 5.4, there exists a bounded open set V in $\mathbb{C}$, a point $\gamma_\infty \in \Gamma^{\mathbb{N}}$, and a number $j \in \mathbb{N}$ such that $\{\gamma_\infty\} \times \partial V \subset \tilde{J}(f) \cap P(f)$, and such that $f_{\gamma,n_j}(y_0) \in V$. Then, we have $\partial V \subset P^*(G)$. Since $g(P^*(G)) \subset P^*(G)$ for each $g \in G$, the maximum principle implies that $V \subset \text{int}(\hat{K}(G))$. Hence, $f_{\gamma,n_j}(y_0) \in \text{int}(\hat{K}(G))$. Therefore, we have proved Lemma 5.6. □

We now demonstrate Theorem 3.10-1 and Theorem 3.10-2.

Proof of Theorem 3.10-1 and Theorem 3.10-2: First, we will show the following claim.
Claim 1. Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$. Then, for any point $y_0 \in F_\gamma(f)$, there exists no non-constant limit function of $\{f_{\gamma,n}\}_{n \in \mathbb{N}}$ around y_0.

To show this claim, suppose that there exists a strictly increasing sequence $\{n_j\}_{j \in \mathbb{N}}$ of positive integers such that f_{γ,n_j} tends to a non-constant map as $j \to \infty$ around y_0. We consider the following two cases: Case (1): $\Gamma \setminus \Gamma_{\min}$ is compact. Case (2): $\Gamma \setminus \Gamma_{\min}$ is not compact. Suppose that we have Case (1). By Lemma 5.6, there exists a number $k \in \mathbb{N}$ such that $f_{\gamma,n_k}(y_0) \in \text{int}(\hat{K}(G))$. Hence, we get that the sequence $\{f_{\sigma^{n_k}(\gamma), n_{k+j} - n_k}\}_{j \in \mathbb{N}}$ converges to a non-constant map around the point $y_1 := f_{\gamma,n_k}(y_0) \in \text{int}(\hat{K}(G))$. However, since we are assuming that $\Gamma \setminus \Gamma_{\min}$ is compact, [27, Theorem 2.20.5(b)] implies that $\cup_{h \in \Gamma \setminus \Gamma_{\min}} h(\hat{K}(G))$ is a compact subset of $\text{int}(\hat{K}(G))$, which implies that if we take the hyperbolic metric for each connected component of $\text{int}(\hat{K}(G))$, then there exists a constant $0 < c < 1$ such that for each $z \in \text{int}(\hat{K}(G))$ and each $h \in \Gamma \setminus \Gamma_{\min}$, we have $\|h'(z)\| \leq c$, where $\|h'(z)\|$ denotes the norm of the derivative of h at z measured from the hyperbolic metric on the connected component W_1 of $\text{int}(\hat{K}(G))$ containing z to that of the connected component W_2 of $\text{int}(\hat{K}(G))$ containing $h(z)$. This causes a contradiction, since we have that $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$ and the sequence $\{f_{\sigma^{n_k}(\gamma), n_{k+j} - n_k}\}_{j \in \mathbb{N}}$ converges to a non-constant map around the point $y_1 \in \text{int}(\hat{K}(G))$. We now suppose that we have Case (2). Then, combining the arguments in Case (1) and [27, Theorem 2.20.5(b), Proposition 2.33], we again obtain a contradiction. Hence, we have shown Claim 1.

Next, let S be a non-empty compact subset of $\Gamma \setminus \Gamma_{\min}$ and let $\gamma \in R(\Gamma, S)$. We show the following claim.
Claim 2. For each point y_0 in each bounded component of $F_\gamma(f)$, there exists a number $n \in \mathbb{N}$ such that $f_{\gamma,n}(y_0) \in \text{int}(\hat{K}(G))$.

To show this claim, suppose that there exists no $n \in \mathbb{N}$ such that $f_{\gamma,n}(y_0) \in \text{int}(\hat{K}(G))$, and we will deduce a contradiction. By Claim 1, $\{f_{\gamma,n}\}_{n \in \mathbb{N}}$ has only constant limit functions around y_0. Moreover, if a point $w_0 \in \mathbb{C}$ is a constant limit function of $\{f_{\gamma,n}\}_{n \in \mathbb{N}}$, then by [28, Lemma 3.13] we must have $w_0 \in P^*(G) \subset \hat{K}(G)$. Since we are assuming that there exists no $n \in \mathbb{N}$ such that $f_{\gamma,n}(y_0) \in \text{int}(\hat{K}(G))$, it follows that $w_0 \in \partial \hat{K}(G)$. Combining it with [27, Theorem 2.20.2] we obtain $w_0 \in \partial \hat{K}(G) \subset J_{\min}$. From this argument, we get that

$$d(f_{\gamma,n}(y_0), J_{\min}) \to 0, \text{ as } n \to \infty. \tag{1}$$

However, since γ belongs to $R(\Gamma, S)$, the above (1) implies that the sequence $\{f_{\gamma,n}(y_0)\}_{n\in\mathbb{N}}$ accumulates in the compact set $\cup_{h\in S}h^{-1}(J_{\min})$, which is apart from $J_{\min}$, by [27, Theorem 2.20.5(b)]. This contradicts (1). Hence, we have shown that Claim 2 holds.

Next, we show the following claim.

Claim 3. There exists exactly one bounded component U_γ of $F_\gamma(f)$.

To show this claim, we take an element $h \in \Gamma_{\min}$ (note that $\Gamma_{\min} \neq \emptyset$, by Proposition 2.10). We write the element γ as $\gamma = (\gamma_1, \gamma_2, \ldots)$. For any $l \in \mathbb{N}$ with $l \geq 2$, let $s_l \in \mathbb{N}$ be an integer with $s_l > l$ such that $\gamma_{s_l} \in S$. We may assume that for each $l \in \mathbb{N}$, $s_l < s_{l+1}$. For each $l \in \mathbb{N}$, let $\gamma^l := (\gamma_1, \gamma_2, \ldots, \gamma_{s_l-1}, h, h, h, \ldots) \in \Gamma^{\mathbb{N}}$ and $\tilde{\gamma}^l := \sigma^{s_l-1}(\gamma) = (\gamma_{s_l}, \gamma_{s_l+1}, \ldots) \in \Gamma^{\mathbb{N}}$. Moreover, let $\rho := (h, h, h, \ldots) \in \Gamma^{\mathbb{N}}$. Since $h \in \Gamma_{\min}$, we have

$$J_\rho(f) = J(h) \subset J_{\min}. \tag{2}$$

Moreover, since γ_{s_l} does not belong to $\Gamma_{\min}$, combining it with [27, Theorem 2.20.5(b)], we obtain $\gamma_{s_l}^{-1}(J(G)) \cap J_{\min} = \emptyset$. Hence, we have that for each $l \in \mathbb{N}$,

$$J_{\tilde{\gamma}^l}(f) = \gamma_{s_l}^{-1}(J_{\sigma^{s_l}(\gamma)}(f)) \subset \gamma_{s_l}^{-1}(J(G)) \subset \hat{\mathbb{C}} \setminus J_{\min}. \tag{3}$$

Combining (2), (3), and [28, Lemma 3.9] we obtain

$$J_\rho(f) < J_{\tilde{\gamma}^l}(f), \tag{4}$$

which implies

$$J_{\gamma^l}(f) = (f_{\gamma,s_l-1})^{-1}(J_\rho(f)) < (f_{\gamma,s_l-1})^{-1}(J_{\tilde{\gamma}^l}(f)) = J_\gamma(f). \tag{5}$$

From [28, Lemma 3.9] and (5), it follows that there exists a bounded component U_γ of $F_\gamma(f)$ such that for each $l \in \mathbb{N}$ with $l \geq 2$,

$$J_{\gamma^l}(f) \subset U_\gamma. \tag{6}$$

We now suppose that there exists a bounded component V of $F_\gamma(f)$ with $V \neq U_\gamma$, and we will deduce a contradiction. Under the above assumption, we take a point $y \in V$. Then, by Claim 2, we get that there exists a number $l \in \mathbb{N}$ such that $f_{\gamma,l}(y) \in \mathrm{int}(\hat{K}(G))$. Since $s_l > l$, we obtain $f_{\gamma,s_l-1}(y) \in \mathrm{int}(\hat{K}(G)) \subset K(h)$, where, $h \in \Gamma_{\min}$ is the element which we have taken before. By (4), we have that there exists a bounded component B of $F_{\tilde{\gamma}^l}(f)$ containing $K(h)$. Hence, we have $f_{\gamma,s_l-1}(y) \in B$. Since the map $f_{\gamma,s_l-1} : V \to B$ is surjective, it follows that $V \cap \big((f_{\gamma,s_l-1})^{-1}(J(h))\big) \neq \emptyset$. Combined with $(f_{\gamma,s_l-1})^{-1}(J(h)) = (f_{\gamma^l,s_l-1})^{-1}(J(h)) = J_{\gamma^l}(f)$, we obtain $V \cap J_{\gamma^l}(f) \neq \emptyset$. However, this causes a contradiction, since we have (6) and $U_\gamma \cap V = \emptyset$. Hence, we have shown Claim 3.

Next, we show the following claim.

Claim 4. $\partial U_\gamma = \partial A_\gamma(f) = J_\gamma(f)$.

To show this claim, since $U_\gamma = \mathrm{int}(K_\gamma(f))$, [28, Lemma 3.4.5] implies that $\partial U_\gamma = J_\gamma(f)$. Moreover, by [28, Lemma 3.4.4] we have $\partial A_\gamma(f) = J_\gamma(f)$. Thus, we have shown Claim 4.

We now show the following claim.

Claim 5. $\hat{J}_\gamma(f) = J_\gamma(f)$ and the map $\omega \mapsto J_\omega(f)$ is continuous at γ with respect to the Hausdorff metric in the space of non-empty compact subsets of $\hat{\mathbb{C}}$.

To show this claim, suppose that there exists a point z with $z \in \hat{J}_\gamma(f) \setminus J_\gamma(f)$. Since $\hat{J}_\gamma(f) \setminus J_\gamma(f)$ is included in the union of bounded components of $F_\gamma(f)$, combining it with Claim 2, we get that there exists a number $n \in \mathbb{N}$ such that $f_{\gamma,n}(z) \in \mathrm{int}(\hat{K}(G)) \subset F(G)$. However, since $z \in \hat{J}_\gamma(f)$, we must have that $f_{\gamma,n}(z) = \pi_{\hat{\mathbb{C}}}(f_\gamma^n(z)) \in \pi_{\hat{\mathbb{C}}}(\tilde{J}(f)) = J(G)$. This is a contradiction. Hence, we obtain $\hat{J}_\gamma(f) = J_\gamma(f)$. Combining it with [28, Lemma 3.4.2], it follows that $\omega \mapsto J_\omega(f)$ is continuous at γ. Therefore, we have shown Claim 5.

Combining all Claims $1, \ldots, 5$, it follows that statements 1, 2a, 2b, and 2c in Theorem 3.10 hold.

We now show statement 2d. Let $\gamma \in R(\Gamma, S)$ be an element. Suppose that $m_2(J_\gamma(f)) > 0$, where m_2 denotes the 2-dimensional Lebesgue measure. Then, there exists a Lebesgue density point $b \in J_\gamma(f)$ so that

$$\lim_{s\to 0} \frac{m_2\left(D(b,s)\cap J_\gamma(f)\right)}{m_2(D(b,s))} = 1. \tag{7}$$

Since γ belongs to $R(\Gamma, S)$, there exists an element $\gamma_\infty \in S$ and a sequence $\{n_j\}_{j\in\mathbb{N}}$ of positive integers such that $n_j \to \infty$ and $\gamma_{n_j} \to \gamma_\infty$ as $j \to \infty$, and such that for each $j \in \mathbb{N}$, $\gamma_{n_j} \in S$. We set $b_j := f_{\gamma,n_j-1}(b)$, for each $j \in \mathbb{N}$. We may assume that there exists a point $a \in \mathbb{C}$ such that $b_j \to a$ as $j \to \infty$. Since $\gamma_{n_j}(b_j) = f_{\gamma,n_j}(b) = \pi_{\hat{\mathbb{C}}}(f_\gamma^{n_j}(\gamma, b)) \in \pi_{\hat{\mathbb{C}}}(\tilde{J}(f)) = J(G)$, we obtain $a \in \gamma_\infty^{-1}(J(G))$. Moreover, by [27, Theorem 2.20.5(b)], we obtain

$$a \in \gamma_\infty^{-1}(J(G)) \subset \mathbb{C} \setminus J_{\min}. \tag{8}$$

Combining it with [27, Theorem 2.20.2], it follows that $r := \inf\{|a-b| \mid b \in P^*(G)\} > 0$. Let ϵ be arbitrary number with $0 < \epsilon < \frac{r}{10}$. We may assume that for each $j \in \mathbb{N}$, we have $b_j \in D(a, \frac{\epsilon}{2})$. For each $j \in \mathbb{N}$, let φ_j be the well-defined inverse branch of $(f_{\gamma,n_j-1})^{-1}$ on $D(a,r)$ such that $\varphi_j(b_j) = b$. Let $V_j := \varphi_j(D(b_j, r-\epsilon))$, for each $j \in \mathbb{N}$. We now show the following claim.
Claim 6. diam $V_j \to 0$, as $\jmath \to \infty$.

To show this claim, suppose that this is not true. Then, there exists a strictly increasing sequence $\{j_k\}_{k\in\mathbb{N}}$ of positive integers and a positive constant κ such that for each $k \in \mathbb{N}$, diam $V_{j_k} \geq \kappa$. From Koebe distortion theorem, it follows that there exists a positive constant c_0 such that for each $k \in \mathbb{N}$, $V_{j_k} \supset D(b, c_0)$. This implies that for each $k \in \mathbb{N}$, $f_{\gamma,v_k}(D(b,c_0)) \subset D(b_{j_k}, r-\epsilon)$, where $v_k := n_{j_k} - 1$. Since $v_k \to \infty$ as $k \to \infty$ and $f_{\gamma',n}|_{F_\infty(G)} \to \infty$ for any $\gamma' \in \Gamma^{\mathbb{N}}$, it follows that for any $n \in \mathbb{N}$, $f_{\gamma,n}(D(b,c_0)) \subset (\hat{\mathbb{C}} \setminus F_\infty(G))$, which implies that $b \in F_\gamma(f)$. However, it contradicts $b \in J_\gamma(f)$. Hence, Claim 6 holds.

Combining Koebe distortion theorem and Claim 6, we see that there exist a constant $K > 0$ and two sequences $\{r_j\}_{j\in\mathbb{N}}$ and $\{R_j\}_{j\in\mathbb{N}}$ of positive numbers such that $K \leq \frac{r_j}{R_j} < 1$ and $D(b, r_j) \subset V_j \subset D(b, R_j)$ for each $j \in \mathbb{N}$, and such that $R_j \to 0$ as $j \to \infty$. From (7), it follows that

$$\lim_{j\to\infty} \frac{m_2\left(V_j \cap F_\gamma(f)\right)}{m_2(V_j)} = 0. \tag{9}$$

For each $j \in \mathbb{N}$, let $\psi_j : D(0,1) \to \varphi_j(D(a,r))$ be a biholomorphic map such that $\psi_j(0) = b$. Then, there exists a constant $0 < c_1 < 1$ such that for each $j \in \mathbb{N}$,

$$\psi_j^{-1}(V_j) \subset D(0, c_1). \tag{10}$$

Combining it with (9) and Koebe distortion theorem, it follows that

$$\lim_{j\to\infty} \frac{m_2\left(\psi_j^{-1}(V_j \cap F_\gamma(f))\right)}{m_2(\psi_j^{-1}(V_j))} = 0. \tag{11}$$

Since $\varphi_j^{-1}(\psi_j(D(0,1)) \subset D(a,r)$ for each $j \in \mathbb{N}$, combining (10) and Cauchy's formula yields that there exists a constant $c_2 > 0$ such that for any $j \in \mathbb{N}$,

$$|(f_{\gamma,n_j-1} \circ \psi_j)'(z)| \leq c_2 \ \text{ on } \psi_j^{-1}(V_j). \tag{12}$$

Combining (11) and (12), we obtain

$$\begin{aligned} \frac{m_2\left(D(b_j, r-\epsilon) \cap F_{\sigma^{n_j-1}(\gamma)}(f)\right)}{m_2(D(b_j, r-\epsilon))} &= \frac{m_2\left((f_{\gamma,n_j-1} \circ \psi_j)(\psi_j^{-1}(V_j \cap F_\gamma(f)))\right)}{m_2(D(b_j, r-\epsilon))} \\ &= \frac{\int_{\psi_j^{-1}(V_j \cap F_\gamma(f))} |(f_{\gamma,n_j-1} \circ \psi_j)'(z)|^2 \, dm_2(z)}{m_2(\psi_j^{-1}(V_j))} \cdot \frac{m_2(\psi_j^{-1}(V_j))}{m_2(D(b_j, r-\epsilon))} \to 0, \end{aligned}$$

as $j \to \infty$. Hence, we obtain

$$\lim_{j\to\infty} \frac{m_2\left(D(b_j, r-\epsilon) \cap J_{\sigma^{n_j-1}(\gamma)}(f)\right)}{m_2(D(b_j, r-\epsilon))} = 1.$$

Since $J_{\sigma^{n_j-1}(\gamma)}(f) \subset J(G)$ for each $j \in \mathbb{N}$, and $b_j \to a$ as $j \to \infty$, it follows that

$$\frac{m_2(D(a, r-\epsilon) \cap J(G))}{m_2(D(a, r-\epsilon))} = 1.$$

This implies that $D(a, r-\epsilon) \subset J(G)$. Since this is valid for any ϵ, we must have that $D(a, r) \subset J(G)$. It follows that the point a belongs to a connected component J of $J(G)$ such that $J \cap P^*(G) \neq \emptyset$. However, [27, Theorem 2.20.2] implies that the component J is equal to $J_{\min}$, which causes a contradiction since we have (8). Hence, we have shown statement 2d in Theorem 3.10-2.

Therefore, we have proved Theorem 3.10-1 and Theorem 3.10-2. □

We now demonstrate Theorem 3.10-3.

Proof of Theorem 3.10-3: First, we remark that the subset $W_{S,p}$ of $\Gamma^{\mathbb{N}}$ is a σ-invariant compact set. Hence, $\overline{f} : W_{S,p} \times \hat{\mathbb{C}} \to W_{S,p} \times \hat{\mathbb{C}}$ is a polynomial skew product over $\sigma : W_{S,p} \to W_{S,p}$. Suppose that $\tilde{J}(\overline{f}) \cap P(\overline{f}) \neq \emptyset$ and let $(\gamma, y) \in \tilde{J}(\overline{f}) \cap P(\overline{f})$ be a point. Then, since the point $\gamma = (\gamma_1, \gamma_2, \dots)$ belongs to $W_{S,p}$, there exists a number $j \in \mathbb{N}$ such that $\gamma_j \in S$. Combining it with [27, Theorem 2.20. 5(b), Theorem 2.20.2] we have $\gamma_j^{-1}(J(G)) \subset \mathbb{C} \setminus \hat{K}(G) \subset \mathbb{C} \setminus P(G)$. Moreover, we have that $\pi_{\hat{\mathbb{C}}}(\overline{f}_\gamma^{j-1}(\gamma, y)) = \pi_{\hat{\mathbb{C}}}(f_\gamma^{j-1}(\gamma, y)) \in J_{\sigma^{j-1}(\gamma)}(f) = \gamma_j^{-1}\left(J_{\sigma^j(\gamma)}(f)\right) \subset \gamma_j^{-1}(J(G))$. Hence, we obtain

$$\pi_{\hat{\mathbb{C}}}(\overline{f}_\gamma^{j-1}(\gamma, y)) \in \mathbb{C} \setminus P(G). \tag{13}$$

However, since $(\gamma, y) \in P(\overline{f})$, we have that $\pi_{\hat{\mathbb{C}}}(\overline{f}_\gamma^{j-1}(\gamma, y)) \in \pi_{\hat{\mathbb{C}}}(P(\overline{f})) \subset P(G)$, which contradicts (13). Hence, we must have that $\tilde{J}(\overline{f}) \cap P(\overline{f}) = \emptyset$. Therefore, $\overline{f} : W_{S,p} \times \hat{\mathbb{C}} \to W_{S,p} \times \hat{\mathbb{C}}$ is a hyperbolic polynomial skew product over the shift map $\sigma : W_{S,p} \to W_{S,p}$.

Combining this with Theorem 3.10-2a and [28, Theorem 4.1] we conclude that there exists a constant $K_{S,p} \geq 1$ such that for each $\gamma \in W_{S,p}$, $J_\gamma(\overline{f})$ is a $K_{S,p}$-quasicircle. Moreover, by Theorem 3.10-2c, we have $J_\gamma(\overline{f}) = J_\gamma(f) = \hat{J}_\gamma(f)$.

Hence, we have shown Theorem 3.10-3. □

We now demonstrate Theorem 3.12.

Proof of Theorem 3.12: Let $\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min})$ and $y \in \operatorname{int}(K_\gamma(f))$. Combining Theorem 3.10-1 and [21, Lemma 1.10], we obtain $\liminf_{n\to\infty} d(f_{\gamma,n}(y),\ J(G)) > 0$. Combining this with [28, Lemma 3.13] and Theorem 3.10-1, we see that there exists a point $a \in P^*(G) \cap F(G)$ such that $\liminf_{n\to\infty} d(f_{\gamma,n}(y), a) = 0$. Since $P^*(G) \cap F(G) \subset \operatorname{int}(\hat{K}(G))$, it follows that there exists a positive integer l such that $f_{\gamma,l}(y) \in \operatorname{int}(\hat{K}(G))$. Combining this and the same method as that in the proof of Claim 3 in the proof of Theorem 3.10-1 and Theorem 3.10-2, we get that there exists exactly one bounded component U_γ of $F_\gamma(f)$. Combining it with [28, Proposition 4.6], it follows that $J_\gamma(f)$ is a Jordan curve. Moreover, by [21, Theorem 2.14-(4)], we have $\hat{J}_\gamma(f) = J_\gamma(f)$.

Thus, we have proved Theorem 3.12. □

We now demonstrate Theorem 3.13.

Proof of Theorem 3.13: Let V be an open set with $J(G) \cap V \neq \emptyset$. We may assume that V is connected. Then, by [10, Corollary 3.1] there exists an element $\alpha_1 \in G$ such that $J(\alpha_1) \cap V \neq \emptyset$. Since we have $G \in \mathcal{G}_{dis}$, [27, Theorem 2.1] implies that there exists an element $\alpha_2 \in G$ such that no connected component J of $J(G)$ satisfies $J(\alpha_1) \cup J(\alpha_2) \subset J$. Hence, we have $\langle \alpha_1, \alpha_2 \rangle \in \mathcal{G}_{dis}$. Since $J(\alpha_1) \cap V \neq \emptyset$, combined with [28, Lemma 3.4.2] we get that there exists an $l_0 \in \mathbb{N}$ such that for each l with $l \geq l_0$, $J(\alpha_2\alpha_1^l) \cap V \neq \emptyset$. Moreover, since no connected component J of $J(G)$ satisfies $J(\alpha_1) \cup J(\alpha_2) \subset J$, [28, Lemma 3.4.2] implies that there exists an $l_1 \in \mathbb{N}$ such that for each l with $l \geq l_1$, $J(\alpha_2\alpha_1^l) \cap J(\alpha_1\alpha_2^l) = \emptyset$. We fix an $l \in \mathbb{N}$ with $l \geq \max\{l_0, l_1\}$. We now show the following claim.

Claim 1. The semigroup $H_0 := \langle \alpha_2\alpha_1^l, \alpha_1\alpha_2^l \rangle$ is hyperbolic, and for the skew product $\tilde{f} : \Gamma_0^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma_0^{\mathbb{N}} \times \hat{\mathbb{C}}$ associated with $\Gamma_0 = \{\alpha_2\alpha_1^l, \alpha_1\alpha_2^l\}$, there exists a constant $K \geq 1$ such that for any $\gamma \in \Gamma_0^{\mathbb{N}}$, $J_\gamma(\tilde{f})$ is a K-quasicircle.

To show this claim, applying Theorem 3.10-3 with $\Gamma = \{\alpha_1, \alpha_2\}$, $S = \Gamma \setminus \Gamma_{\min}$, and $p = 2l+1$, we see that the polynomial skew product $\overline{f} : W_{S,2l+1} \times \hat{\mathbb{C}} \to W_{S,2l+1} \times \hat{\mathbb{C}}$ over $\sigma : W_{S,2l+1} \to W_{S,2l+1}$ is hyperbolic, and that there exists a constant $K \geq 1$ such that for each $\gamma \in W_{S,2l+1}$, $J_\gamma(\overline{f})$ is a K-quasicircle. Moreover, combining the hyperbolicity of $\overline{f}$ above and Remark 5.3, we see that the semigroup H_1 generated by the family $\{\alpha_{j_1} \circ \cdots \circ \alpha_{j_{l+1}} \mid 1 \leq \exists k_1 \leq l+1 \text{ with } j_{k_1} = 1, 1 \leq \exists k_2 \leq l+1 \text{ with } j_{k_2} = 2\}$ is hyperbolic. Hence, the semigroup H_0, which is a subsemigroup of H_1, is hyperbolic. Therefore, Claim 1 holds.

We now show the following claim.

Claim 2. We have either $J(\alpha_2\alpha_1^l) < J(\alpha_1\alpha_2^l)$, or $J(\alpha_1\alpha_2^l) < J(\alpha_2\alpha_1^l)$.

To show this claim, since $J(\alpha_2\alpha_1^l) \cap J(\alpha_1\alpha_2^l) = \emptyset$ and $H_0 \in \mathcal{G}$, combined with [28, Lemma 3.9], we obtain Claim 2.

By Claim 2, we have the following two cases.

Case 1. $J(\alpha_2\alpha_1^l) < J(\alpha_1\alpha_2^l)$.

Case 2. $J(\alpha_1\alpha_2^l) < J(\alpha_2\alpha_1^l)$.

We may assume that we have Case 1 (when we have Case 2, we can show all statements of our theorem, using the same method as below). Let $A := K(\alpha_1\alpha_2^l) \setminus \operatorname{int}(K(\alpha_2\alpha_1^l))$. By Claim 1, we have that $J(\alpha_1\alpha_2^l)$ and $J(\alpha_2\alpha_1^l)$ are quasicircles. Moreover, since $H_0 \in \mathcal{G}_{dis}$ and H_0 is hyperbolic, we must have $P^*(H_0) \subset \operatorname{int}(K(\alpha_2\alpha_1^l))$. Therefore, it follows that if we take a small open neighborhood U of A, then there exists a number $n \in \mathbb{N}$ such that, setting $h_1 := (\alpha_2\alpha_1^l)^n$ and $h_2 := (\alpha_1\alpha_2^l)^n$, we have that

$$h_1^{-1}(\overline{U}) \cup h_2^{-1}(\overline{U}) \subset U \text{ and } h_1^{-1}(\overline{U}) \cap h_2^{-1}(\overline{U}) = \emptyset. \tag{14}$$

Moreover, combining [28, Lemma 3.4.2] and that $J(h_1) \cap V \neq \emptyset$, we get that there exists a $k \in \mathbb{N}$ such that $J(h_2h_1^k) \cap V \neq \emptyset$. We set $g_1 := h_1^{k+1}$ and $g_2 := h_2h_1^k$. Moreover, we set $H := \langle g_1, g_2 \rangle$. Since H is a subsemigroup of H_0 and H_0 is hyperbolic, we have that H is hyperbolic. Moreover, (14) implies that $g_1^{-1}(\overline{U}) \cup g_2^{-1}(\overline{U}) \subset U$ and $g_1^{-1}(\overline{U}) \cap g_2^{-1}(\overline{U}) = \emptyset$. Hence, we have shown that for the semigroup $H = \langle g_1, g_2 \rangle$, statements 1,2, and 3 in Theorem 3.13 hold.

From statement 2 and [10, Corollary 3.2], we obtain $J(H) \subset \overline{U}$ and $g_1^{-1}(J(H)) \cap g_2^{-1}(J(H)) = \emptyset$. Combining this with [20, Lemma 2.4] and [28, Lemma 3.5.2], it follows that the skew product $f : \Gamma_1^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma_1^{\mathbb{N}} \times \hat{\mathbb{C}}$ associated with $\Gamma_1 = \{g_1, g_2\}$ satisfies that $J(H)$ is equal to the disjoint union of the sets $\{\hat{J}_\gamma(f)\}_{\gamma \in \Gamma_1^{\mathbb{N}}}$. Moreover, since H is hyperbolic, [21, Theorem 2.14-(2)] implies that for each $\gamma \in \Gamma_1^{\mathbb{N}}$, $\hat{J}_\gamma(f) = J_\gamma(f)$. In particular, the map $\gamma \mapsto J_\gamma(f)$ from $\Gamma_1^{\mathbb{N}}$ into the space of non-empty compact sets in $\hat{\mathbb{C}}$, is injective. Since $J_\gamma(f)$ is connected for each $\gamma \in \Gamma_1^{\mathbb{N}}$ (Claim 1), it follows that for each connected component J of $J(H)$, there exists an element $\gamma \in \Gamma_1^{\mathbb{N}}$ such that $J = J_\gamma(f)$. Furthermore, by Claim 1, each connected component J of $J(H)$ is a K-quasicircle, where K is a constant not depending on J. Moreover, by [21, Theorem 2.14-(4)], the map $\gamma \mapsto J_\gamma(f)$ from $\Gamma_1^{\mathbb{N}}$ into the space of non-empty compact sets in $\hat{\mathbb{C}}$, is continuous with respect to the Hausdorff metric. Therefore, we have shown that statements 4a,4b,4c, and 4d hold for $H = \langle g_1, g_2 \rangle$ and $f : \Gamma_1^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma_1^{\mathbb{N}} \times \hat{\mathbb{C}}$.

We now show that statement 4e holds. Since we are assuming Case 1, Proposition 2.10 implies that $\{h_1, h_2\}_{\min} = \{h_1\}$. Hence $J(g_1) < J(g_2)$. Combining it with Proposition 2.10 and statement 4b, we obtain

$$J(g_1) = J_{\min}(H) \text{ and } J(g_2) = J_{\max}(H). \tag{15}$$

Moreover, since $J(g_1) = J(\alpha_2\alpha_1^l)$, $J(\alpha_2\alpha_1^l) \cap V \neq \emptyset$, $J(g_2) = J(h_2h_1^k)$, and $J(h_2h_1^k) \cap V \neq \emptyset$, it follows that

$$J_{\min}(H) \cap V \neq \emptyset \text{ and } J_{\max}(H) \cap V \neq \emptyset. \tag{16}$$

Let $\gamma \in \Gamma^{\mathbb{N}}$ be an element such that $J_\gamma(f) \cap (J_{\min}(H) \cup J_{\max}(H)) = \emptyset$. By statement 4b, we obtain

$$J_{\min}(H) < J_\gamma(f) < J_{\max}(H). \tag{17}$$

Since we are assuming V is connected, combining (16) and (17), we obtain $J_\gamma(f) \cap V \neq \emptyset$. Therefore, we have proved that statement 4e holds.

We now show that statement 4f holds. To show that, let $\omega = (\omega_1, \omega_2, \ldots) \in \Gamma_1^{\mathbb{N}}$ be an element such that $\sharp(\{j \in \mathbb{N} \mid \omega_j = g_1\}) = \sharp(\{j \in \mathbb{N} \mid \omega_j = g_2\}) = \infty$. For each $r \in \mathbb{N}$, let $\omega^r = (\omega_{r,1}, \omega_{r,2}, \ldots) \in \Gamma_1^{\mathbb{N}}$ be the element such that $\begin{cases} \omega_{r,j} = \omega_j & (1 \leq j \leq r), \\ \omega_{r,j} = g_1 & (j \geq r+1). \end{cases}$ Moreover, let $\rho^r = (\rho_{r,1}, \rho_{r,2}, \ldots) \in \Gamma_1^{\mathbb{N}}$ be the element such that $\begin{cases} \rho_{r,j} = \omega_j & (1 \leq j \leq r), \\ \rho_{r,j} = g_2 & (j \geq r+1). \end{cases}$ Combining (15), statement 4a, and statement 4b, we see that for each $r \in \mathbb{N}$, $J(g_1) < J_{\sigma^r(\omega)}(f) < J(g_2)$. Hence, by Theorem 2.8-3, we get that for each $r \in \mathbb{N}$, $(f_{\omega,r})^{-1}(J(g_1)) < (f_{\omega,r})^{-1}\left(J_{\sigma^r(\omega)}(f)\right) < (f_{\omega,r})^{-1}(J(g_2))$. Since we have $(f_{\omega,r})^{-1}(J(g_1)) = J_{\omega^r}(f)$, $(f_{\omega,r})^{-1}\left(J_{\sigma^r(\omega)}(f)\right) = J_\omega(f)$, and $(f_{\omega,r})^{-1}(J(g_2)) = J_{\rho^r}(f)$, it follows that

$$J_{\omega^r}(f) < J_\omega(f) < J_{\rho^r}(f), \tag{18}$$

for each $r \in \mathbb{N}$. Moreover, since $\omega^r \to \omega$ and $\rho^r \to \omega$ in $\Gamma_1^{\mathbb{N}}$ as $r \to \infty$, statement 4d implies that $J_{\omega^r}(f) \to J_\omega(f)$ and $J_{\rho^r}(f) \to J_\omega(f)$ as $r \to \infty$, with respect to the Hausdorff metric. Combined with (18), statement 4b and statement 4c, we get that for any connected component W of $F(H)$, we must have $\partial W \cap J_\omega(f) = \emptyset$. Since $F(G) \subset F(H)$, it follows that for any connected component W' of $F(G)$, $\partial W' \cap J_\omega(f) = \emptyset$. Therefore, we have shown that statement 4f holds.

Thus, we have proved Theorem 3.13. □

5.2 Proofs of results in 3.2

In this section, we demonstrate Theorem 3.17. We need the following notations and lemmas.

Definition 5.7. Let h be a polynomial with $\deg(h) \geq 2$. Suppose that $J(h)$ is connected. Let ψ be a biholomorphic map $\hat{\mathbb{C}} \setminus \overline{D(0,1)} \to F_\infty(h)$ with $\psi(\infty) = \infty$ such that $\psi^{-1} \circ h \circ \psi(z) = z^{\deg(h)}$, for each $z \in \hat{\mathbb{C}} \setminus \overline{D(0,1)}$. (For the existence of the biholomorphic map ψ, see [13, Theorem 9.5].) For each $\theta \in \partial D(0,1)$, we set $T(\theta) := \psi(\{r\theta \mid 1 < r \leq \infty\})$. This is called the external ray (for $K(h)$) with angle θ.

Lemma 5.8. *Let h be a polynomial with $\deg(h) \geq 2$. Suppose that $J(h)$ is connected and locally connected and $J(h)$ is not a Jordan curve. Moreover, suppose that there exists an attracting periodic point of h in $K(h)$. Then, for any $\epsilon > 0$, there exist a point $p \in J(h)$ and elements $\theta_1, \theta_2 \in \partial D(0,1)$ with $\theta_1 \neq \theta_2$, such that all of the following hold.*

1. *For each $i = 1, 2$, the external ray $T(\theta_i)$ lands at the point p.*
2. *Let V_1 and V_2 be the two connected components of $\hat{\mathbb{C}} \setminus (T(\theta_1) \cup T(\theta_2) \cup \{p\})$. Then, for each $i = 1, 2$, $V_i \cap J(h) \neq \emptyset$. Moreover, there exists an i such that diam $(V_i \cap K(h)) \leq \epsilon$.*

Proof. Let $\psi : \hat{\mathbb{C}} \setminus \overline{D(0,1)} \to F_\infty(h)$ be a biholomorphic map with $\psi(\infty) = \infty$ such that for each $z \in \hat{\mathbb{C}} \setminus \partial D(0,1), \psi^{-1} \circ h \circ \psi(z) = z^{\deg(h)}$. Since $J(h)$ is locally connected, the map $\psi : \hat{\mathbb{C}} \setminus \overline{D(0,1)} \to F_\infty(h)$ extends continuously over $\partial D(0,1)$, mapping $\partial D(0,1)$ onto $J(h)$. Moreover, since $J(h)$ is not a Jordan curve, it follows that there exist a point $p_0 \in J(h)$ and two points $t_{0,1}, t_{0,2} \in \partial D(0,1)$ with $t_{0,1} \neq t_{0,2}$ such that two external rays $T(t_{0,1})$ and $T(t_{0,2})$ land at the same point p_0. Considering a higher iterate of h if necessary, we may assume that there exists an attracting fixed point of h in $\text{int}(K(h))$. Let $a \in \text{int}(K(h))$ be an attracting fixed point of h

and let U be the connected component of $\mathrm{int}(K(h))$ containing a. Then, there exists a critical point $c \in U$ of h. Let V_0 be the connected component of $\hat{\mathbb{C}} \setminus (T(t_1) \cup T(t_2) \cup \{p_0\})$ containing a. Moreover, for each $n \in \mathbb{N}$, let V_n be the connected component of $(h^n)^{-1}(V_0)$ containing a. Since $c \in U$, we get that for each $n \in \mathbb{N}$, $c \in V_n$. Hence, setting $e_n := \deg(h^n : V_n \to V_0)$, it follows that $e_n \to \infty$ as $n \to \infty$. We fix an $n \in \mathbb{N}$ satisfying $e_n > d$, where $d := \deg(h)$. Since $\deg(h^n : V_n \cap F_\infty(h) \to V_0 \cap F_\infty(h)) = \deg(h^n : V_n \to V_0)$, we have that the number of connected components of $V_n \cap F_\infty(h)$ is equal to e_n. Moreover, every connected component of $V_n \cap F_\infty(h)$ is a connected component of $(h^n)^{-1}(V_0 \cap F_\infty(h))$. Hence, it follows that there exist mutually disjoint arcs $\xi_1, \xi_2, \ldots, \xi_{e_n}$ in $\mathbb{C}$ satisfying all of the following.

1. For each j, $h^n(\xi_j) = (T(t_1) \cup T(t_2) \cup \{p_0\}) \cap \mathbb{C}$.
2. For each j, $\xi_j \cup \{\infty\}$ is the closure of union of two external rays and $\xi_j \cup \{\infty\}$ is a Jordan curve.
3. $\partial V_n = \xi_1 \cup \cdots \cup \xi_{e_n} \cup \{\infty\}$.

For each $j = 1, \ldots, e_n$, let W_j be the connected component of $\hat{\mathbb{C}} \setminus (\xi_j \cup \{\infty\})$ that does not contain V_n. Then, each W_j is a connected component of $\hat{\mathbb{C}} \setminus \overline{V_n}$. Hence, for each (i,j) with $i \neq j$, $W_i \cap W_j = \emptyset$. Since the number of critical values of h in $\mathbb{C}$ is less than or equal to $d-1$, we have that $\sharp(\{1 \leq j \leq e_n \mid W_j \cap CV(h) = \emptyset\}) \geq e_n - (d-1)$. Therefore, denoting by $u_{1,j}$ the number of well-defined inverse branches of h^{-1} on W_j, we obtain $\sum_{j=1}^{e_n} u_{1,j} \geq d(e_n - (d-1)) \geq d$. Inductively, denoting by $u_{k,j}$ the number of well-defined inverse branches of $(h^k)^{-1}$ on W_j, we obtain

$$\sum_{j=1}^{e_n} u_{k,j} \geq d(d-(d-1)) \geq d, \text{for each } k \in \mathbb{N}. \tag{19}$$

For each $k \in \mathbb{N}$, we take a well-defined inverse branch ζ_k of $(h^k)^{-1}$ on a domain W_j, and let $B_k := \zeta_k(W_j)$. Then, $h^k : B_k \to W_j$ is biholomorphic. Since ∂B_k is the closure of finite union of external rays and h^{n+k} maps each connected component of $(\partial B_k) \cap \mathbb{C}$ onto $(T(t_1) \cup T(t_2) \cup \{p_0\}) \cap \mathbb{C}$, B_k is a Jordan domain. Hence, $h^k : B_k \to W_j$ induces a homeomorphism $\partial B_k \cong \partial W_j$. Therefore, ∂B_k is the closure of union of two external rays, which implies that $B_k \cap F_\infty(h)$ is a connected component of $(h^k)^{-1}(W_j \cap F_\infty(h))$. Hence, we obtain

$$l\left(\overline{\psi^{-1}(B_k \cap F_\infty(h))} \cap \partial D(0,1)\right) \to 0 \text{ as } k \to \infty, \tag{20}$$

where $l(\cdot)$ denotes the arc length of a subarc of $\partial D(0,1)$. Since $\psi : \hat{\mathbb{C}} \setminus \overline{D(0,1)} \to F_\infty(h)$ extends continuously over $\partial D(0,1)$, (20) implies that diam $(B_k \cap J(h)) \to 0$ as $k \to \infty$. Hence, there exists a $k \in \mathbb{N}$ such that diam $(B_k \cap K(h)) \leq \epsilon$. Let $\theta_1, \theta_2 \in \partial D(0,1)$ be two elements such that $\partial B_k = \overline{T(\theta_1) \cup T(\theta_2)}$. Then, there exists a point $p \in J(h)$ such that each $T(\theta_i)$ lands at the point p. By [13, Lemma 17.5], any of two connected components of $\hat{\mathbb{C}} \setminus (T(\theta_1) \cup T(\theta_2) \cup \{p\})$ intersects $J(h)$.

Thus, we have proved Lemma 5.8. □

Lemma 5.9. *Let G be a polynomial semigroup generated by a compact subset Γ of $\mathrm{Poly}_{\deg \geq 2}$. Let $f : \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}} \to \Gamma^{\mathbb{N}} \times \hat{\mathbb{C}}$ be the skew product associated with the family Γ. Suppose $G \in \mathcal{G}_{dis}$. Let $m \in \mathbb{N}$ and suppose that there exists an element $(h_1, \ldots, h_m) \in \Gamma^m$ such that setting $h = h_m \circ \cdots \circ h_1$, $J(h)$ is connected and locally connected, and $J(h)$ is not a Jordan curve. Moreover, suppose that there exists an attracting periodic point of h in $K(h)$. Let $\alpha = (\alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$ be the element such that for each $k, l \in \mathbb{N} \cup \{0\}$ with $1 \leq l \leq m$, $\alpha_{km+l} = h_l$. Let $\rho_0 \in \Gamma \setminus \Gamma_{\min}$ be an element and let $\beta = (\rho_0, \alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$. Moreover, let $\psi_\beta : \hat{\mathbb{C}} \setminus \overline{D(0,1)} \to A_\beta(f)$ be a biholomorphic map with $\psi_\beta(\infty) = \infty$. Furthermore, for each $\theta \in \partial D(0,1)$, let $T_\beta(\theta) = \psi_\beta(\{r\theta \mid 1 < r \leq \infty\})$. Then, for any $\epsilon > 0$, there exist a point $p \in J_\beta(f)$ and elements $\theta_1, \theta_2 \in \partial D(0,1)$ with $\theta_1 \neq \theta_2$, such that both of the following statements 1 and 2 hold.*

1. *For each $i = 1, 2$, $T_\beta(\theta_i)$ lands at p.*
2. *Let V_1 and V_2 be the two connected components of $\hat{\mathbb{C}} \setminus (T_\beta(\theta_1) \cup T_\beta(\theta_2) \cup \{p\})$. Then, for each $i = 1, 2$, $V_i \cap J_\beta(f) \neq \emptyset$. Moreover, there exists an i such that diam $(V_i \cap K_\beta(f)) \leq \epsilon$ and such that $V_i \cap J_\beta(f) \subset \rho_0^{-1}(J(G)) \subset \mathbb{C} \setminus P(G)$.*

Proof. We use the notation and argument in the proof of Lemma 5.8. Taking a higher iterate of h, we may assume that $d := \deg(h) > \deg(\rho_0)$. Then, from (19), it follows that for each $k \in \mathbb{N}$, we can take a well-defined inverse branch ζ_k of $(h^k)^{-1}$ on a domain W_j such that setting $B_k := \zeta_k(W_j)$, B_k does not contain any critical value of ρ_0. By (20), there exists a $k \in \mathbb{N}$ such that diam $(B_k \cap J(h)) \leq \epsilon'$, where $\epsilon' > 0$ is a small number. Let B be a connected component of $\rho_0^{-1}(B_k)$. Then, there exist a point $p \in J_\beta(f)$ and elements $\theta_1, \theta_2 \in \partial D(0, 1)$ with $\theta_1 \neq \theta_2$ such that for each $i = 1, 2$, $T_\beta(\theta_i)$ lands at p, and such that B is a connected component of $\hat{\mathbb{C}} \setminus (T_\beta(\theta_1) \cup T_\beta(\theta_2) \cup \{p\})$. Taking ϵ' so small, we obtain diam $(B \cap K_\beta(f)) =$ diam $(B \cap J_\beta(f)) \leq \epsilon$. Moreover, since $\rho_0 \in \Gamma \setminus \Gamma_{\min}$, by [27, Theorem 2.20.5(b), Theorem 2.20.2] we obtain $J_\beta(f) = \rho_0^{-1}(J(h)) \subset \rho_0^{-1}(J(G)) \subset \mathbb{C} \setminus P(G)$. Hence, $B \cap J_\beta(f) \subset \rho_0^{-1}(J(G)) \subset \mathbb{C} \setminus P(G)$. Therefore, we have proved Lemma 5.9. □

We now demonstrate Theorem 3.17-1.

Proof of Theorem 3.17-1: Let γ be as in Theorem 3.17-1. Then, by Theorem 3.12, $J_\gamma(f)$ is a Jordan curve. Moreover, setting $h = h_m \circ \cdots \circ h_1$, since h is hyperbolic and $J(h)$ is not a quasicircle, $J(h)$ is not a Jordan curve. Combining it with [28, Lemma 4.5] and Lemma 5.8, it follows that $J_\gamma(f)$ is not a quasicircle. Moreover, $A_\gamma(f)$ is a John domain (cf. [23, Theorem 1.12]). Combining the above arguments with [14, Theorem 9.3], we conclude that the bounded component U_γ of $F_\gamma(f)$ is not a John domain.

Thus, we have proved Theorem 3.17-1. □

We now demonstrate Theorem 3.17-2.

Proof of Theorem 3.17-2: Let ρ_0, β, γ be as in Theorem 3.17-2. By Theorem 3.12, $J_\gamma(f)$ is a Jordan curve. By Theorem 2.9-4, we have $\emptyset \neq \text{int}(\hat{K}(G)) \subset \text{int}(K(h))$. Moreover, h is semi-hyperbolic. Hence, h has an attracting periodic point in $K(h)$. Combining [28, Lemma 4.5] and Lemma 5.9, we get that $J_\gamma(f)$ is not a quasicircle. Combining it with the argument in the proof of Theorem 3.17-1, it follows that $A_\gamma(f)$ is a John domain, but the bounded component U_γ of $F_\gamma(f)$ is not a John domain.

Thus, we have proved Theorem 3.17-2. □

We now prove Theorem 3.18.

Proof of Theorem 3.18: By [26, Theorem 3.17], we have $h_1^{-1}(J(G)) \cap h_2^{-1}(J(G)) = \emptyset$. Combining this with [28, Lemma 3.5.2, Lemma 3.6] and [21, Theorem 2.14(2)], we obtain that $J(G) = \amalg_{\gamma \in \Gamma^{\mathbb{N}}} J_\gamma(f)$ (disjoint union) and for each connected component J of $J(G)$, there exists a unique $\gamma \in \Gamma^{\mathbb{N}}$ such that $J = J_\gamma(f)$.

In order to prove that we have exactly one of statements 1 and 2, suppose that $J(h_1)$ and $J(h_2)$ are Jordan curves. By Proposition 2.10, we may assume that $J_{\min}(G) = J(h_1)$ and $J_{\max}(G) = J(h_2)$. Then by [27, Theorem 2.20.5(b)], we have $\text{int}(K(h_1)) = \text{int}(\hat{K}(G))$. Thus $P^*(G)$ is included in a connected component of $\text{int}(\hat{K}(G))$. Combining it with [28, Proposition 2.25], we obtain that statement 1 in Theorem 3.18 holds.

We now suppose that $J(h_j)$ is not a Jordan curve. Then, as above, we have $J_{\min}(G) = J(h_j)$ and $J_{\max}(G) = J(h_i)$, where $i \neq j$. Then, combining Theorem 3.10-3, Theorem 3.17-1 and [28, Lemma 4.4], we obtain that exactly one of statements (a),(b),(c) in Theorem 3.18-2. Thus, we have proved Theorem 3.18. □

5.3 Proofs of results in 3.3

In this subsection, we will demonstrate results in Section 3.3.

we now prove Corollary 3.20.

Proof of Corollary 3.20: By Remark 3.7, there exists a compact subset S of $\Gamma \setminus \Gamma_{\min}$ such that the interior of S with respect to the space Γ is not empty. Let $\mathcal{U} := R(\Gamma, S)$. Then, it is easy to see that $\mathcal{U}$ is residual in $\Gamma^{\mathbb{N}}$, and that for each Borel probability measure τ on $\text{Poly}_{\deg \geq 2}$ with $\Gamma_\tau = \Gamma$, we have $\tilde{\tau}(\mathcal{U}) = 1$. Moreover, by Theorem 3.10-1 and Theorem 3.10-2, each $\gamma \in \mathcal{U}$ satisfies properties 1,2,3, and 4 in Corollary 3.20. Hence, we have proved Corollary 3.20. $\square$

We now prove Corollary 3.21.

Proof of Corollary 3.21: Setting $\mathcal{U} := \{\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min}) \mid \exists \{n_k\} \text{ s.t. } \sigma^{n_k}(\gamma) \to \alpha\}$, it is easy to see that statement 1 holds. We now prove statement 2. From our assumption, there exist $h_1, \ldots, h_m \in \Gamma$ such that $J(h_m \circ \cdots \circ h_1)$ is not a quasicircle. Let $\alpha = (\alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$ be such that for each $k, l \in \mathbb{N} \cup \{0\}$ with $1 \leq l \leq m$, $\alpha_{km+l} = h_l$. Let $\rho_0 \in \Gamma \setminus \Gamma_{\min}$ be an element and let $\beta = (\rho_0, \alpha_1, \alpha_2, \ldots) \in \Gamma^{\mathbb{N}}$. Let $\mathcal{V} := \{\gamma \in R(\Gamma, \Gamma \setminus \Gamma_{\min}) \mid \exists \{n_k\} \text{ s.t. } \sigma^{n_k}(\gamma) \to \beta\}$. Then $\mathcal{V}$ satisfies the desired properties. $\square$

References

[1] R. Brück, *Geometric properties of Julia sets of the composition of polynomials of the form* $z^2 + c_n$, Pacific J. Math., **198** (2001), no. 2, 347–372.

[2] R. Brück, M. Büger and S. Reitz, *Random iterations of polynomials of the form* $z^2 + c_n$*: Connectedness of Julia sets*, Ergodic Theory Dynam. Systems, **19**, (1999), No.5, 1221–1231.

[3] M. Büger, *Self-similarity of Julia sets of the composition of polynomials*, Ergodic Theory Dynam. Systems, **17** (1997), 1289–1297.

[4] M. Büger, *On the composition of polynomials of the form* $z^2 + c_n$, Math. Ann. 310 (1998), no. 4, 661–683.

[5] L. Carleson, P. W. Jones and J. -C. Yoccoz, *Julia and John,* Bol. Soc. Bras. Mat. **25, N.1** 1994, 1–30.

[6] R. Devaney, *An Introduction to Chaotic Dynamical Systems*, Perseus Books, 1989.

[7] J. E. Fornaess and N. Sibony, *Random iterations of rational functions*, Ergodic Theory Dynam. Systems, **11**(1991), 687–708.

[8] Z. Gong, W. Qiu and Y. Li, *Connectedness of Julia sets for a quadratic random dynamical system*, Ergodic Theory Dynam. Systems, (2003), **23**, 1807-1815.

[9] Z. Gong and F. Ren, *A random dynamical system formed by infinitely many functions,* Journal of Fudan University, **35**, 1996, 387–392.

[10] A. Hinkkanen and G. J. Martin, *The Dynamics of Semigroups of Rational Functions I*, Proc. London Math. Soc. (3)**73**(1996), 358–384.

[11] M. Jonsson, *Ergodic properties of fibered rational maps* , Ark. Mat., 38 (2000), pp 281–317.

[12] O. Lehto and K. I. Virtanen, *Quasiconformal Mappings in the plane,* Springer-Verlag, 1973.

[13] J. Milnor, *Dynamics in One Complex Variable* (Third Edition), Annals of Mathematical Studies, Number 160, Princeton University Press, 2006.

[14] R. Näkki and J. Väisälä, *John Discs,* Expo. math. 9(1991), 3–43.

[15] K. Pilgrim and Tan Lei, *Rational maps with disconnected Julia set*, Asterisque 261, (2000), 349–384.

[16] R. Stankewitz, T. Sugawa, and H. Sumi, *Some counterexamples in dynamics of rational semigroups*, Annales Academiae Scientiarum Fennicae Mathematica Vol. 29, 2004, 357–366.

[17] R. Stankewitz and H. Sumi, *Dynamical properties and structure of Julia sets of postcritically bounded polynomial semigroups*, to appear in Trans. Amer. Math. Soc., http://arxiv.org/abs/0708.3187.

[18] D. Steinsaltz, *Random logistic maps Lyapnov exponents*, Indag. Mathem., N. S., 12 (4), 557–584, 2001.

[19] H. Sumi, *On Dynamics of Hyperbolic Rational Semigroups,* Journal of Mathematics of Kyoto University, Vol. 37, No. 4, 1997, 717–733.

[20] H. Sumi, *Skew product maps related to finitely generated rational semigroups,* Nonlinearity, **13**, (2000), 995–1019.

[21] H. Sumi, *Dynamics of sub-hyperbolic and semi-hyperbolic rational semigroups and skew products*, Ergodic Theory Dynam. Systems, (2001), **21**, 563–603.

[22] H. Sumi, *Dimensions of Julia sets of expanding rational semigroups*, Kodai Mathematical Journal, Vol. 28, No. 2, 2005, pp390–422. (See also http://arxiv.org/abs/math.DS/0405522.)

[23] H. Sumi, *Semi-hyperbolic fibered rational maps and rational semigroups*, Ergodic Theory Dynam. Systems (2006), **26**, 893–922.

[24] H. Sumi, *The space of postcritically bounded 2-generator polynomial semigroups with hyperbolicity*, RIMS Kokyuroku 1494, 62–86, 2006. (Proceeding paper.)

[25] H. Sumi, *Random dynamics of polynomials and devil's-staircase-like functions in the complex plane*, Applied Mathematics and Computation 187 (2007) pp489-500. (Proceeding paper.)

[26] H. Sumi, *Interaction cohomology of forward or backward self-similar systems*, Adv. Math., 222 (2009), no. 3, 729–781.

[27] H. Sumi, *Dynamics of postcritically bounded polynomial semigroups I: connected components of the Julia sets*, to appear in Discrete and Continuous Dynamical Systems - Ser. A, http://arxiv.org/abs/0811.3664.

[28] H. Sumi, *Dynamics of postcritically bounded polynomial semigroups III: classification of semi-hyperbolic semigroups and random Julia sets which are Jordan curves but not quasicircles*, to appear in Ergodic Theory Dynam. Systems, http://arxiv.org/abs/0811.4536.

[29] H. Sumi, *Random complex dynamics and semigroups of holomorphic maps*, to appear in Proc. London Math. Soc., http://arxiv.org/abs/0812.4483.

[30] H. Sumi, *Rational semigroups, random complex dynamics and singular functions on the complex plane*, survey article, to appear in Sugaku Expositions.

[31] H. Sumi, *Cooperation principle in random complex dynamics and singular functions on the complex plane*, to appear in RIMS Kokyuroku. (Proceedings paper.)

[32] H. Sumi, in preparation.

[33] H. Sumi and M. Urbanski, *Real analyticity of Hausdorff dimension for expanding rational semigroups*, Ergodic Theory Dynam. Systems (2010), Vol. 30, No. 2, 601-633.

[34] H. Sumi and M. Urbanski, *The equilibrium states for semigroups of rational maps*, Monatsh. Math., 156 (2009), no. 4, 371–390.

[35] Y. Sun and C-C. Yang. *On the connectivity of the Julia set of a finitely generated rational semigroup*, Proc. Amer. Math. Soc., Vol. 130, No. 1, 49–52, 2001.